




NEUROPATHIC PAIN: AMYGDALOID STRESS AS CULPRIT AND CORTICAL STIMULATION AS TREATMENT

BORISS SAGALAJEV



NEUROPATHIC PAIN: AMYGDALOID STRESS AS CULPRIT AND CORTICAL STIMULATION AS TREATMENT

Boriss Sagalajev

Department of Physiology
Faculty of Medicine
University of Helsinki
Finland

Doctoral Programme in Drug Research

ACADEMIC DISSERTATION

To be presented, with the permission of the Faculty of Medicine, University of Helsinki, for public examination in Lecture Hall 2, Biomedicum 1, Haartmaninkatu 8, on August 4, 2017, at 12:00 noon.

Helsinki 2017

Supervisor **Antti Pertovaara**, MD, PhD

Professor of Physiology
Department of Physiology
Faculty of Medicine
University of Helsinki
Helsinki, Finland

Reviewers **Raimo Tuominen**, MD, PhD

Professor of Pharmacology and Toxicology
Division of Pharmacology and Pharmacotherapy
Faculty of Pharmacy
University of Helsinki
Helsinki, Finland

Turo Nurmikko, MD, PhD

Professor in Pain Relief
Neuroscience Research Centre
The Walton Centre NHS Foundation Trust

Hon. Professor of Pain Science
Faculty of Health and Life Sciences
Clinical Sciences Centre
University of Liverpool
Liverpool, UK

Opponent **Rashid Giniatullin**, MD, PhD, DMSc

Professor of Cell Biology
Department of Neurobiology
A.I. Virtanen Institute for Molecular Sciences
University of Eastern Finland
Kuopio, Finland

ISBN 978-951-51-3575-9 (paperback)

ISBN 978-951-51-3576-6 (PDF)

Unigrafia

Helsinki 2017

To teachers, friends, and family!

CONTENTS

ABBREVIATIONS.....	1
EPIGRAPH	3
ABSTRACT	4
POPULAR ABSTRACT IN FINNISH	5
LIST OF ORIGINAL PUBLICATIONS.....	7
1 LITERATURE REVIEW	9
1.1 Introduction	9
1.2 Nociceptive pathways.....	10
1.2.1 Primary afferent fibers and their spinal targets.....	10
1.2.2 Gate control theory of pain.....	11
1.2.3 Ascending nociceptive pathways.....	13
1.2.4 Descending nociceptive pathways.....	13
1.3 Medullospinal neurons.....	15
1.4 Amygdala	16
1.4.1 Role of the amygdala in Pavlovian conditioning	16
1.4.2 Role of the amygdala in nociception	16
1.4.3 Chronic pain-induced neuroplastic changes in the amygdala	18
1.4.4 Role of the amygdala in stress.....	18
1.4.4.1 General adaptation syndrome	18
1.4.4.2 Hypothalamic-pituitary-adrenocortical axis	19
1.4.4.3 Corticotropin-releasing factor.....	19
1.4.4.4 Reactive oxygen species	20
1.4.4.5 Transient receptor potential channels	22
1.5 Somatosensory cortex.....	22
2 AIMS	24
3 METHODS	25
3.1 Ethical considerations.....	25
3.2 Drugs.....	26
3.3 Surgeries	26

3.3.1 Anesthesia	26
3.3.2 Spared nerve injury model	27
3.3.3 Spinal nerve ligation model	28
3.3.4 Intracranial cannula implantation	29
3.3.5 Intrathecal catheterization	29
3.4 Behavioral tests	29
3.4.1 Mechanical pain	29
3.4.2 Heat pain	29
3.4.3 Emotional pain	29
3.5 Electrophysiology	30
4 RESULTS AND DISCUSSION	31
4.1 Medullospinal 5-HT neurons as a relay for bidirectional modulation of pain hypersensitivity by the amygdala (I, IV)	31
4.2 Reactive oxygen species as a culprit for pronociceptive hyperactivity of the amygdala (II, IV)	35
4.3 Medullospinal 5-HT neurons as a relay for attenuation of neuropathic pain by cortical stimulation (III)	38
4.4 General discussion	43
5 CAVEATS	45
6 FUTURE PROSPECTS	46
7 CONCLUSIONS	48
8 REFERENCES	50

ABBREVIATIONS

5-HT	serotonin (5-hydroxytryptamine)
8-OH-DPAT	8-hydroxy-2-(dipropylamino)tetralin
BLA	basolateral amygdaloid nucleus
CeA	central amygdaloid nucleus
Cg	cingulate cortex
CHEM	CHEMBRIDGE-5861528
CT	central thalamus
DMH	dorsomedial hypothalamus
DRT	dorsal reticular nucleus (caudal)
GABA	γ -aminobutyric acid
GLU	glutamate
I	insular cortex
ITC	intercalated cell mass
LA	lateral amygdaloid nucleus
LC	locus coeruleus
LH	lateral hypothalamus
LIDO	lidocaine
M1	primary motor cortex
MDT	medial dorsal thalamus
mGLUR	metabotropic glutamate receptor
mPFC	medial prefrontal cortex
NMDA	N-methyl-D-aspartate
PAG	periaqueductal gray
PbN	parabrachial nucleus
PBN	phenyl-N-tert-butyl nitron
PVN	paraventricular nucleus
ROS	reactive oxygen species
RtF	reticular formation
RVM	rostral ventromedial medulla
S1	primary somatosensory cortex
S2	secondary somatosensory cortex
SC	superior colliculus
SDH	spinal dorsal horn
SNI	spared nerve injury
SNL	spinal nerve ligation
Str	striatum
t-BOOH	tert-butyl-hydroperoxide
TEMPOL	4-hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl

ThCN	thalamocortical network
TMS	transcranial magnetic stimulation
TRP	transient receptor potential
TRPA	ankyrin subfamily of TRP channels
TRPC	canonical subfamily of TRP channels
VLM	ventrolateral medulla (caudal)
VPT	ventral posterior thalamus
WDR	wide-dynamic range

EPIGRAPH

“It was in this way that the gray cub learned other attributes of his mother than the soft, soothing tongue. In his insistent crawling toward the light, he discovered in her a nose that with a sharp nudge administered rebuke, and later, a paw, that crushed him down or rolled him over and over with swift, calculating stroke. Thus he learned hurt; and on top of it he learned to avoid hurt, first, by not incurring the risk of it; and second, when he had incurred the risk, by dodging and by retreating. These were conscious actions, and were the results of his first generalizations upon the world. Before that he had recoiled automatically from hurt, as he had crawled automatically toward the light. After that he recoiled from hurt because he *knew* that it was hurt.”

White Fang

Jack London

1906

ABSTRACT

Despite extensive research on the mechanisms of nociceptive pain, little is known about the processes that lead to neuropathic pain development. As one of the most severe and drug-resistant forms of chronic pain, neuropathic pain represents a major burden for patients, their families, and society. Therefore, new insights into the pathophysiology of neuropathic pain are needed.

The amygdala represents a complex of brain nuclei responsible for mediating negative emotions, such as fear. Moreover, maladaptive changes in the amygdala contribute to the development of chronic pain and its comorbidities, such as anxiety and depression. However, little is known about the mechanisms for amygdala-mediated pain hypersensitivity in neuropathy. We therefore investigated descending pathways engaged by the amygdala for modulation of spinal nociception in neuropathic rats.

In particular, we investigated the contribution of oxidative stress in the amygdala on the development of neuropathic pain. Oxidative stress is characterized by overproduction and poor detoxification of reactive oxygen species (ROS). ROS act as endogenous agonists of transient receptor potential (TRP) channels, which account for transduction of noxious stimulation to a nociceptive signal in peripheral nerves and for nociceptive transmission in the spinal cord. We therefore assessed how blocking TRP channels and detoxification of ROS with antioxidants in the amygdala influence pain hypersensitivity and pain affect in neuropathic rats.

Furthermore, we investigated whether electrical cortical stimulation can reverse changes associated with pronociceptive activity of the amygdala. In particular, we examined descending pathways recruited by stimulation of the secondary somatosensory cortex (S2), which has proven to be an efficacious site for attenuation of drug-resistant neuropathic pain by transcranial magnetic stimulation (TMS) in patients.

In the present series of studies, we demonstrate that N-methyl-D-aspartate (NMDA), TRPA₁, and TRPC_{4/5} channels in the amygdala contribute to the development of neuropathic pain and that amygdaloid treatment with antioxidants can attenuate this pain. Furthermore, we provide evidence that S2 stimulation suppresses spinal nociception in neuropathic animals with hypersensitivity, but not in animals without hypersensitivity or in control animals. Medullospinal serotonergic pathways acting on the spinal 5-HT_{1A} receptor underlie the descending pain inhibition in neuropathic animals both following treatment of the amygdala with antioxidants or TRPA₁ antagonists and following S2 cortex stimulation.

POPULAR ABSTRACT IN FINNISH

Hermovauriokipu: mantelitumakkeen hapetusstressi aiheuttajana ja aivokuoren stimulaatio hoitokeinona

Vaikka kivun fysiologisia mekanismeja on tutkittu runsaasti, niin hermovaurion aiheuttaman neuropaattisen kivun ja kivulle herkistymisen mekanismit ovat vielä monilta osin tuntemattomia. Hermovauriokipu on usein kroonista eli pitkäkestoista ja nykyisin käytössä olevat hoitomuodot tehoavat hermovauriokipuun usein huonosti. Yleisyytensä ja vaikeahoitoisuutensa vuoksi krooninen hermovauriokipu on merkittävä hyvinvointia heikentävä tekijä yhteiskunnassa. Siksi onkin tarpeen saada lisätietoa hermovauriokivun patofysiologisista mekanismeista, joiden parempi tuntemus voi auttaa kehittämään tehokkaampia hoitomuotoja.

Mantelitumake on keskeinen aivorakenne negatiivisille emootioille kuten pelolle. Patofysiologisissa tiloissa syntyvien mantelitumakkeen rakenteellisten ja toiminnallisten muutosten tiedetään myötävaikuttavan kroonisten kiputilojen sekä niihin liittyvien emotionaalisten häiriötilojen kuten ahdistuksen ja pelon syntyyn. Mekanismit, joilla mantelitumake voi myötävaikuttaa kivulle herkistymiseen hermovauriotilanteissa, ovat vielä huonosti tunnettuja. Tässä työsarjassa tutkittiin mantelitumakkeesta lähtevän ja selkäyttimeen laskevan kivun säätelyn mekanismeja kokeellisessa hermovauriokivun rottamallissa.

Erityisesti tutkimuksen kohteena oli mantelitumakkeen hapetusstressin merkitys hermovauriokivun kehittymiselle. Hapetusstressiin tyypillisesti kuuluu hapetusstressituotteiden ylituotanto sekä vähentynyt hajotus. Hapetusstressituotteet voivat tunnetusti aktivoida hermosolun kalvon TRP-ionikanavia, joiden toiminta aikaansaa ja voimistaa kipuviestejä. Tässä työsarjassa selvitettiin, kuinka mantelitumakkeeseen paikallisesti annostellut TRP-ionikanavien salpaajat tai hapetusstressituotteiden vastavaikuttajat vaikuttivat hermovaurion aiheuttamaan kivulle herkistymiseen ja kivun affektiivisiin vasteeseen.

Tämän lisäksi työsarjassa selvitettiin voiko aivokuorelle annetulla sähköstimulaatiolla vaimentaa hermovauriokipua. Erityisesti tutkittiin sekundaarisen (S2) tuntoaivokuoren stimulaation vaikutusta ja mekanismeja, koska aiemmissa ihmistöissä on S2-aivokuoren transkraniaalisella magneettistimulaatiolla aikaansaatua tehokas kivun lievitys potilailla, jotka ovat kärsineet hermovauriokivusta, johon tavanomainen lääkehoito on tehonnut huonosti.

Tämän työsarjan tulokset osoittavat, että mantelitumakkeen N-metyyli-D-aspartaatti reseptorit sekä TRPA₁ ja TRPC_{4/5} ionikavat myötävaikuttavat

hermovammaan liittyvän kivulle herkistymistilan kehittymiseen. Myös mantelitulmakkeen hapetusstressi, jonka synnyttämät tuotteet ovat TRP-ionikanavia aktivoivia tekijöitä, on tulosten mukaan myötävaikuttamassa kivulle herkistymiseen hermovaurion jälkeen. S2-aivokuoren sähköstimulaatio vaimensi eläinten kipuviestejä selkäydintasolla, mutta vain jos eläimellä hermovaurio ja siihen liittyvä herkistyminen kosketusärsykkeille. S2-aivokuoren sähköstimulaation samoin kuin mantelitulmakkeen TRP-kanavien salpauksen ja hapetusstressin eston hermovauriokipua lievittävä vaikutus välittyi selkäydintasolle ydinjatkoksesta selkäytimeen laskevien serotonergisten solujen välityksellä, jotka selkäytimen takasarveen serotoniinia vapauttaessaan aktivoivat 5-HT_{1A} reseptoreita, jotka vaimentavat kipuviestejä selkäytimen tasolla.

LIST OF ORIGINAL PUBLICATIONS

- I **Sagalajev B**, Bourbia N, Beloushko E, Wei H, Pertovaara A. Bidirectional amygdaloid control of neuropathic hypersensitivity mediated by descending serotonergic pathways acting on spinal 5-HT₃ and 5-HT_{1A} receptors. *Behav Brain Res* 2015;282:14-24. doi: 10.1016/j.bbr.2014.12.052.
- II Wei H, **Sagalajev B***, Yüzer MA, Koivisto A, Pertovaara A. Regulation of neuropathic pain behavior by amygdaloid TRPC4/C5 channels. *Neurosci Lett* 2015;608:12-7. doi: 10.1016/j.neulet.2015.09.033.
- III **Sagalajev B**, Viisanen H, Wei H, Pertovaara A. Descending antinociception induced by secondary somatosensory cortex stimulation in experimental neuropathy: role of the medullospinal serotonergic pathway. *J Neurophysiol* 2017;117(3):1200-1214. doi: 10.1152/jn.00836.2016.
- IV **Sagalajev B**, Wei H, Li Z, Albayrak I, Chen Z, Tian L, Koivisto A, Pertovaara A. Oxidative stress and TRPA1 in the amygdala contribute to neuropathic pain by suppressing descending pathways acting on the 5-HT_{1A} receptor. Submitted.

* Equal contribution

1 LITERATURE REVIEW

1.1 Introduction

Actual or potential tissue injury results from noxious stimulation and leads to acquisition of an unpleasant sensory experience, namely pain. Pain serves as an alarm system that leads to development of acute withdrawal and chronic aversive behavior for minimization of further encounter with noxious stimulation. Accordingly, pain raises our chances for a longer and healthier life. Individuals with a rare congenital insensitivity to pain usually die from severe injuries in early adulthood. This is when parents are likely to stop monitoring these individuals for tissue damage and compensating for pain malfunction [28]. At the opposite extreme are patients who experience pain from innocuous stimulation (i.e., allodynia), increased pain from noxious stimulation (i.e., hyperalgesia), or both (i.e., hypersensitivity). The lives of such patients is full of limitations and, therefore, can result in development of depression and, eventually, suicidal behavior. Although pain is subjective, it is always unpleasant. Therefore, experiences (e.g. pricking), that comprise only the sensorial aspect of pain, but fail to elicit negative emotions, are not recognized as pain by the International Association for the Study of Pain [109]. This is consistent with reports of lobotomized patients, in whom destruction of the cortex responsible for encoding of pain unpleasantness kept pain sensations from becoming bothersome [130]. At the same time, experiences, that comprise only the emotional aspect of pain [42], can be regarded as pain. In fact, such experiences can lead to development of somatic (e.g. headache) [15,53] or visceral painful sensations (e.g. chest pain) [34,37,76].

Typically, however, pain arises from activation of pain receptors, or nociceptors. Pain that results from normal somatosensory system functioning is called nociceptive, whereas pain that results from abnormal somatosensory functioning is called neuropathic. Peripheral neuropathic pain develops in the setting of peripheral nerve injury, which triggers pronociceptive changes in primary afferent fibers. In addition, pronociceptive changes can take place at the spinal and supraspinal levels and, in fact, outlast changes in the periphery [20]. In the short term, central pronociceptive changes are only functional. In the long term, these changes can lead to structural alterations [137,138]. Neuropathic pain represents the most severe and drug-resistant form of chronic pain. In the clinic, less than a half of neuropathic patients receive satisfactory pain treatment. These numbers are even lower outside the clinic. One of the main reasons for this lies in the necessity for progressive increase in the drug dosage that unavoidably leads to development of intolerable side effects. Furthermore, neuropathic pain predicts adverse

consequences for the patients' families and a substantial economic burden on society in the form of healthcare costs and patients' inability to work [119,120].

Although the processes leading to chronic pain development remain largely unknown, some of its steps are noteworthy. Patients with chronic pain display an increase in neuronal sodium channel expression [36] that likely accounts for development of hyperexcitability and spontaneous firing in nociceptors. Similarly, animals with experimental persistent pain display greater expression of cation channels, hyperexcitability of nociceptors, and activation of previously silent nociceptors [35,73]. Spinal mechanisms for facilitation of nociceptive transmission involve long-term potentiation (LTP) of synaptic strength between active nociceptive afferents and target neurons [84]. Spinal LTP requires proinflammatory activation of microglia, which may also account for a decrease in the inhibitory potency of a γ -aminobutyric acid (GABA) [26,27]. Furthermore, in the setting of nerve injury, non-nociceptive afferents can display a nociceptive phenotype [117] and, together with nociceptive afferents, sprout new branches in the spinal dorsal horn (SDH) [182] (however, see [7,63]). Altogether, this amplifies nociceptive input to the brain that, in the long-term, can lead to maladaptive changes, such as a decrease in descending inhibition and an increase in descending facilitation of ascending nociceptive transmission (for a review, see [57]).

1.2 Nociceptive pathways

1.2.1 Primary afferent fibers and their spinal targets

In primary afferent nerve fibers, conduction velocity decreases with the diameter of the fiber and thickness of myelin sheath in the following order: proprioceptive A α , mechanosensitive A β , thermosensitive A δ , thermosensitive C (unmyelinated). In addition, A δ - and C-fibers can respond to polymodal noxious stimulation and, therefore, are responsible for nociception (for a review, see [105]). Nociceptive information from A δ -fibers is first to arrive at the spinal cord and thus to the brain for perception of sharp and well-localized pain. Information from C-fibers arrives 1 to 2 seconds later and results in perception of burning and poorly localized pain, which increases in intensity with repetition of noxious stimulation. Furthermore, this second pain can outlast noxious stimulation and activation of C-fibers, and, therefore, likely, plays a greater role in neuropathic pain than A δ -fibers [127].

Primary afferents project to the SDH, from where they follow separate pathways. A δ - and C-fibers form synapses with interneurons as well as with projection neurons, axons of which cross the midline and ascend through the anterolateral column to the brain. In addition, non-nociceptive fibers of primary afferents form

synapses in the SDH, although their major branches ascend to the brain directly through the ipsilateral dorsal column. In the superficial layers of the SDH, projection neurons are mostly nociceptive specific, whereas, in deeper layers, they act as wide-dynamic-range (WDR) neurons, which converge nociceptive and non-nociceptive inputs [175].

1.2.2 Gate control theory of pain

Although reciprocal principles for encoding of nociceptive and non-nociceptive information in the spinal cord remain largely unknown, some of them appear to be consistent with the gate control theory of pain [108]. The theory rests on observations that, although both non-nociceptive and nociceptive afferents excite projection neurons, they display opposite effects on spontaneous inhibition of projection neurons by interneurons. In particular, non-nociceptive afferents directly facilitate inhibitory action of interneurons, whereas nociceptive afferents indirectly attenuate it (Fig. 1). The theory therefore suggests a circuitry, in which branches of non-nociceptive afferents can reduce ascending nociceptive transmission by “closing” the gates to projection neurons.

Interestingly, the gate control theory also indicates that, because non-nociceptive information is first to arrive at the somatosensory cortex, it likely sets receptivity of cortical neurons for the following nociceptive input and activates descending pathways for modulation of the gate control system in the spinal cord. Moreover, because the somatosensory cortex displays somatotopic arrangement, it likely triggers descending modulation capable of achieving a high level of spatial accuracy.

In addition, the gate control theory suggests that, because projection neurons receive nociceptive information both from visceral and somatic afferents (viscerosomatic convergence), the former may “open” the gates for the latter and, thus, refer the sense of pain from the viscera to elsewhere in the body. The theory therefore explains why myocardial ischemia may lead to pain development, for instance, in the left shoulder.




Figure 1. Gate control theory of pain. Projection neurons in the spinal dorsal horn receive main excitatory input from nociceptive C-fibers and minor excitatory input from non-nociceptive Aβ-fibers. In addition, C-fibers can excite a chain of two inhibitory interneurons and, thereby, disinhibit nociceptive transmission through projection neurons. Aβ-fibers, on the other hand, can impede such disinhibition by exciting the second inhibitory interneuron.

1.2.3 Ascending nociceptive pathways

In the brain (Fig. 2), projection neurons form the most prominent connections with the thalamus, to which they ascend directly or via the brainstem reticular formation [181]. The thalamus, in turn, relays nociceptive information to the somatosensory cortex, which encodes the sensory-discriminative aspect of pain, and to the anterior-cingulate [174] and insular cortices [21], which, in contrast, encode the affective-motivational aspect. Additional projection neurons ascend to the superior colliculus (SC), which promotes visual identification of noxious stimulation [131], to the periaqueductal gray (PAG), which acts as the opioidergic center for pain modulation [124], and to the parabrachial nucleus (PbN), which constitutes a part of the ascending arousal system [150]. Furthermore, the PbN relays nociceptive information to the hypothalamus [102,104] for regulation of endocrine responses and to the central nucleus of the amygdala (CeA) for development of avoidance behavior [12,90].

1.2.4 Descending nociceptive pathways

Medullary neurons descend to the spinal cord (Fig. 2), where they modulate nociceptive transmission from primary afferent fibers to projection neurons [56,93,151,158]. In turn, the rostral ventromedial part of the medulla (RVM) represents a target for direct modulation by higher brain structures, such as the locus coeruleus, PbN, PAG, SC, and hypothalamus [19,22,139,157,167,176]. Furthermore, indirect sources of RVM modulation can originate from the CeA, striatum (Str), primary motor cortex and likely from the S2 [4,9,10,49,59,86,87,171-173]. Such circuitry forms the neural basis for appropriate adjustment of pain threshold in a constantly changing environment. Chronic nerve injury, however, can lead to its dysregulation and, thereby, to development of pain hypersensitivity.




Figure 2. Ascending (left) and descending (right) nociceptive pathways.

CeA	central amygdaloid nucleus	PbN	parabrachial nucleus
Cg	cingulate cortex	PVN	paraventricular nucleus
CT	central thalamus	RtF	reticular formation
DMH	dorsomedial hypothalamus	RVM	rostral ventromedial medulla
DRt	dorsal reticular nucleus (caudal)	S1	primary somatosensory cortex
I	insular cortex	S2	secondary somatosensory cortex
LC	locus coeruleus	SC	superior colliculus
LH	lateral hypothalamus	Str	striatum
M1	primary motor cortex	VLM	ventrolateral medulla (caudal)
MDT	medial dorsal thalamus	VPT	ventral posterior thalamus
PAG	periaqueductal gray		

1.3 Medullospinal neurons

The RVM projects to the spinal cord with axonal collaterals terminating along the rostrocaudal axis in both the spinal ventral horn and SDH [93]. It thereby acts as a relay for descending modulation of the vasomotor tone and nociception [33]. Originating from the RVM serotonergic (5-hydroxytryptamine; 5-HT) medullospinal neurons display partial topographic specificity. In particular, neurons that display co-localization of 5-HT with peptides terminate in the ventral horn and deep layers of the SDH [61,75,107]. On the other hand, neurons that display co-localization of 5-HT with GABA terminate exclusively in superficial layers of the SDH [24,107].

GABA acts as an inhibitory neurotransmitter in healthy animals, and, as evident at the supraspinal level, 5-HT neurons can express GABA receptors [170]. Therefore, GABA may calibrate the release of 5-HT from medullospinal neurons and its diffusion to adjacent neurons (i.e., volume transmission [1]) by the means of autoinhibition in the SDH. At a high level of volume transmission, all neurons can act as a 5-HT target. Depending on the receptor type they express, however, activation of a 5-HT receptor can result in either inhibition (e.g. 5-HT₁) or facilitation (e.g. 5-HT₂₋₄) of neuronal activity (for a review, see [41]). Furthermore, whether such modulation of neuronal activity in the SDH leads to pro- or antinociception depends on the role 5-HT target neurons play in nociception (transmission, facilitation, or inhibition).

Activation of GABA_A receptors leads to an influx of chloride anions and cell hyperpolarization. The transmembrane gradient of chloride anions is then restored by the chloride exporter KCC2. In neuropathic conditions, however, a decrease in KCC2 expression leads to a lower inhibitory potency of GABA. Such loss of GABA-mediated inhibition in the SDH contributes to development of chronic pain [27], whereas in the ventral horn, this loss leads to development of spasticity. In the latter case, however, a decrease in KCC2 expression results from the loss of descending 5-HT input [18]. It is therefore likely that chronic pain develops (at least in part) for a similar reason [32,46].

In the RVM, 5-HT neurons display slow and steady discharge that fails to change in response to noxious stimulation. Such discharge is exclusive to 5-HT neurons and indicates their role in tonic modulation of pain [106]. For dynamic modulation, on the other hand, non-5-HT neurons are most likely responsible. Non-5-HT neurons respond to noxious stimulation with either transient inhibition (OFF cells) or facilitation (ON cells) of their discharge. In OFF cells, ongoing discharge positively correlates with antinociception, whereas in ON cells, ongoing discharge positively correlates with pronociception [56]. The role of non-5-HT neurons that fail to respond to noxious stimulation, however, remains unclear. Together with 5-HT

neurons, unresponsive to noxious stimulation non-5-HT neurons constitute the population of NEUTRAL cells [106].

1.4 Amygdala

1.4.1 Role of the amygdala in Pavlovian conditioning

After pairing with an unpleasant stimulus, a neutral cue can become aversive, whereas, after pairing with a pleasant stimulus, it can become appetitive. Thus, depending on the conditions, a neutral cue can acquire either a negative or a positive emotional value. Such conditioning was first demonstrated more than a century ago by Ivan Pavlov, who could induce salivation in dogs with a sound of a bell after pairing it with feeding. Furthermore, replacement of an auditory with a noxious signal produced the same effect without any signs of pain. A reward of feeding likely compensated for the unpleasantness of noxious stimulation and, in a sense, led to the dogs' comprehension of the phrase "no pain, no gain". However, noxious stimulation of a body part, which remained intact during pairing, did result in unbearable pain [108,122]. In Pavlovian conditioning, pain inhibition therefore requires a circuitry for precise localization of noxious stimulation and previous experience. Later studies revealed that this circuitry incorporates a limbic brain area, the amygdala, which can indirectly (via the RVM) regulate processing of nociception throughout the spinal cord and, thereby, modulate pain threshold at different body sites [58,59,93]. In addition, the amygdala acts as a junction of pathways for polymodal sensations and emotions. The amygdala predetermines whether collision of nociceptive and non-nociceptive signals will lead to the ability of the latter to elicit negative emotions in the future, such as fear [90]. Pain-related fear represents acute stress that can raise the pain threshold and prepare the animal for the fight-or-flight response [58,59]. In chronic stress, however, fear transitions to anxiety and susceptibility to pain increases [91]. Chronic stress and anxiety represent major complaints among neuropathic patients (e.g. [129,132]) and imply that the amygdala exerts a role in pain chronification [72,74,94,112,147].

1.4.2 Role of the amygdala in nociception

The amygdala (Fig. 3) consists of a group of nuclei, each having distinct connections with pain-modulating brain structures (for a review, see [115]). The lateral amygdaloid nucleus (LA) receives polymodal sensory information from the thalamocortical network and sends it to the basolateral amygdaloid nucleus (BLA), which, in turn, relays it to the central amygdaloid nucleus (CeA) and to the medial

prefrontal cortex (mPFC). The BLA projects to the CeA either directly or, as the mPFC, via intercalated mass of inhibitory interneurons (ITC cells) in the amygdala [71,143]. The CeA, in addition, receives nociceptive-specific information from the spinal cord via the PbN and pairs it with inhibitory inputs from ITC cells or with direct modulatory inputs from the BLA [12,102]. The outcome of such pairing is then transmitted through the PAG to the RVM [59], which, as the CeA, also receives nociceptive-specific information from the PbN [139]. The RVM compares information from the PAG with information from the PbN for regulation of nociceptive processing in the SDH [59].




Figure 3. Role of the amygdala in nociception.

BLA	basolateral amygdaloid nucleus	- ->	amygdaloid nociceptive circuitry
CeA	central amygdaloid nucleus	->	ascending nociceptive pathways
ITC	intercalated cell mass	=>	descending nociceptive pathways
LA	lateral amygdaloid nucleus		
mPFC	medial prefrontal cortex		
PAG	periaqueductal gray		
PbN	parabrachial nucleus		
RVM	rostral ventromedial medulla		
SDH	spinal dorsal horn		
ThCN	thalamocortical network		

1.4.3 Chronic pain-induced neuroplastic changes in the amygdala

Projections of the BLA to the mPFC allow pain to affect cognitive functions, whereas projections of the mPFC to amygdaloid ITC cells permit cognition to modulate pain. In chronic pain states, however, the BLA displays hyperactivity that leads to preferential excitation of inhibitory interneurons rather than of pyramidal cells in the mPFC. A decrease in capability of the mPFC to excite inhibitory ITC cells in the amygdala results in disinhibition of the CeA. Taken together, this leads to impairment of cognitive functions and pain persistence [71]. The reason for hyperactivity of amygdaloid nuclei, however, does not lie in the consistent afferent input from chronic injury. Rather, the injury leads to pain maintenance by changing the strength of amygdaloid synapses. In brain slices of animals with chronic pain, potentiation of synaptic transmission from LA to BLA [71], from BLA to CeA [44,116], as well as from PbN to CeA [65] has been shown. Brain imaging studies revealed amygdaloid hyperactivity in arthritic patients [85], whereas neuropathic patients display a decrease in amygdaloid activity as a result of long- but not short-term pain relief [47]. GABAergic inhibition of the amygdala in neuropathic animals leads to attenuation of pain hypersensitivity and affect [123], whereas stimulation with corticosterone in healthy animals leads to development of behavioral and spinal hypersensitivity [51,128]. Overall, these findings support the hypothesis that peripheral nerve injury increases amygdaloid activity, which accounts for supraspinal pain chronification.

1.4.4 Role of the amygdala in stress

1.4.4.1 General adaptation syndrome

In biology, interchange between damage and defense induces stress, which consists of three stages. The first stage features activation of the sympathetic nervous system for execution of defense behavior or, in other words, of the fight-or-flight response. The second stage involves the synthesis and release of glucocorticoids into the bloodstream to help to cope with stress. However, if stress continues, the third stage takes place, in which exhaustion, malfunction, and degradation of the body are evident. The first and second stages imply a normal general adaptation to stress agents (stressors), whereas the third stage indicates emergence of the adaptation disorder. Moreover, our susceptibility to stressors as well as their combination and specificity of action determine what type of a disorder develops [148]. In this light, neuropathic pain—a disorder in its own right—represents a result of abnormal adaptation to nerve injury. Therefore, expanding our

understanding of the mechanisms of stress maladaptations in the setting of nerve injury can provide new insights into the basis for neuropathic pain.

1.4.4.2 Hypothalamic-pituitary-adrenocortical axis

Acute stress activates sympathetic system, which enhances brain activity through elevation of sugar and oxygen blood levels. This allows rapid decision-making on how to minimize external danger. In addition, stress triggers a slower, yet a longer-lasting response characterized by a release of corticotropin-releasing factor (CRF) from the paraventricular nucleus (PVN) of the hypothalamus (for reviews, see [16,60]). CRF stimulates production of adrenocorticotrophic hormone (ACTH) and beta-endorphin from pro-opiomelanocortin in the pituitary gland [103]. ACTH promotes production of glucocorticoids in the adrenal cortex, whereas beta-endorphin suppresses pain. The latter, in part, explains the analgesia evident in acute stress, during which, however, non-opioid mechanisms also take place [145].

Experimental lesions of the BLA and medial amygdaloid nucleus in rats reveal a decrease in activation of the hypothalamic-pituitary-adrenocortical (HPA) axis during acute emotional stress [13,29], whereas lesions of the CeA fail to have any effect [29,126]. On the other hand, lesions of the CeA dampen responses of the HPA axis to acute inflammatory stress [185] and late activation of the parasympathetic system in acute pain stress. Furthermore, CeA lesions abolish both early sympathetic and late parasympathetic responses to stimuli paired with acute pain stress [140]. Taken together, these studies demonstrate that amygdaloid nuclei play a pivotal role in stress regulation by initiating stress responses of the HPA axis and recovery from stress-induced sympathetic activation, as well as by forming and replaying stress memory (i.e., conditioning).

1.4.4.3 Corticotropin-releasing factor

Interestingly, the amygdala represents a major site for expression of CRF and its receptors outside the hypothalamus, suggesting a leading role of CRF circuitry in orchestrating stress and pain responses. In particular, the lateral hypothalamus (LH, through a release of CRF) and PbN (through a release of calcitonin gene-related peptide) innervate CRF-containing neurons in the CeA. In return, CRF-containing neurons in the CeA project back to the LH and PbN. It is likely that these CRF projections from the CeA regulate ascending nociceptive transmission through the LH and PbN. In addition, the CeA both directly and indirectly (through the bed nucleus of the stria terminalis) innervates CRF-containing neurons in the PVN of the

hypothalamus and, thus, likely regulates stress responses of the HPA axis (for a review, see [50]).

CRF in the amygdala acts on two G-protein-coupled receptors: an excitatory CRF₁ receptor and inhibitory CRF₂ receptor [69]. Blocking CRF₁ receptors in the CeA prevents shock-induced freezing [6], whereas its activation leads to synaptic facilitation between the CeA and parabrachial terminals and to greater neuronal output of the CeA [67]. In addition, chronic pain leads to synaptic facilitation between the CeA and the parabrachial terminals as well as between the CeA and BLA. However, blocking CRF₁ receptors can inhibit synaptic facilitation, whereas blocking CRF₂ receptors can facilitate it [45]. CRF₁ receptors require lower doses of CRF for CeA stimulation than CRF₂ receptors do for CeA inhibition [69]. Together, amygdaloid CRF₁ and CRF₂ receptors are likely to be involved in pro- and antinociceptive actions driven by the amygdala.

Interestingly, although rats with high stress reactivity display pronociception mediated by CRF₁ receptors [66], activation of CRF₁ receptors in normal rats results in pronociception, independent of the HPA axis [67]. The amygdala therefore likely regulates pain through CRF circuitry distinct from the HPA axis. Repetitive activation of the HPA axis, however, can initiate pronociceptive changes in amygdaloid expression of CRF and its receptors. This is consistent with findings showing that prolonged exposure of the CeA to glucocorticoids raises CRF expression levels in the CeA of healthy rats that leads to pain facilitation and, likely through sensitization of the HPA axis, to development of anxiety [51,149,162]. Neuropathic animals also fail to display any functional deficiencies in the HPA axis and have higher CRF expression levels in the CeA [141,166].

1.4.4.4 Reactive oxygen species

Glucocorticoids help maintain high brain activity levels by facilitating glucose availability through glycogenolysis, gluconeogenesis, and lipolysis. Moreover, glucocorticoids suppress the immune system that results in even greater availability of glucose for brain consumption (for reviews, see [125,154,155]). In chronic stress, however, suppression of immune responses and an increase in brain metabolism raise the risk of cell damage from uncontrolled ROS production (for a review, see [152]). Animal models of stress, among which are those driven by noxious stimulation, reveal a dynamic correlation between brain antioxidative activity and behavior. Initiation of antioxidative activity in the first week of stress corresponds to a decrease in exploratory behavior, which most likely indicates development of fear. In the second week of stress, antioxidative activity reaches its peak and behavior returns to normal. Beyond this period, however, antioxidative activity fails

and animals display abnormally high exploratory behavior [52], which may mask development of anxiety [156]. The amygdala therefore, as an essential brain area for development of fear and anxiety, is likely to be susceptible to oxidative changes in long-term stress. Moreover, since nerve injury can represent a form of long-term stress, similar changes in the amygdala likely take place during transition of pain from an acute to a chronic state.

Induction of hyperalgesia and comorbid anxiety with peripheral administration of pain-inducing agents is associated with an increase in amygdaloid levels of ROS, which not only promotes excitatory transmission from the PbN to the CeA but also elevates excitability and activity of the CeA [68,101]. Similarly, activation of a metabotropic glutamate receptor type 5 (mGLUR5) in the amygdala induces hyperalgesia in healthy rats by promoting ROS production and, thereby, elevating CeA activity [97]. The table below demonstrates that mGLUR5 is likely the only metabotropic glutamate receptor, which CeA neurons express on their cell bodies (Table 1). In light of this notion and the fact that the CeA acts as a sole nucleus of the amygdala that projects directly to regions, which either transmit ascending nociception or modulate it, mGLUR5 emerges as the main culprit for amygdala-driven pronociception [3,14,54,95,96,133,134].

Table 1. Glutamate receptors in the amygdala and effect of their activation.

Family	Group	Type	Synapse	Localization	CeA activity
Metabotropic	I	mGLUR1	ITC*→CeA	presynaptic	↑
		mGLUR5	BLA→CeA	postsynaptic	↑
	II	mGLUR2,3	ITC*→CeA	presynaptic	↓
	III	mGLUR4,7,8	PbN→CeA	presynaptic	↓
		mGLUR7	ITC*→CeA	presynaptic	↑
		mGLUR8	BLA→CeA	presynaptic	↓
Ionotropic		NMDA	PbN→CeA	postsynaptic	↑

BLA basolateral amygdaloid nucleus * indirect evidence
 CeA central amygdaloid nucleus
 ITC intercalated cell mass
 PbN parabrachial nucleus

1.4.4.5 Transient receptor potential channels

Transient receptor potential (TRP) channels are non-selective cation channels that are highly sensitive to cellular redox status. Increase in ROS production leads to TRP-mediated influx of cations, which can depolarize the cell membrane and act as second messengers (for a review, see [121]).

From the ankyrin subfamily of TRP channels, mammalian cells express only type 1 (TRPA₁), which acts as a sensor for irritant chemicals at C-fibers in the skin [77] and lungs [114]. There, activation of TRPA₁ channels by ROS can lead to sensitization of C-fibers [2,144] and, thus, to development of pain hypersensitivity [31]. Furthermore, in the SDH of the spinal cord, presynaptic TRPA₁ channels facilitate glutamatergic transmission from primary afferents [80] that likely accounts for development of secondary hypersensitivity, as evident both in patients [83] and experimental animals [177]. Similarly, presynaptic TRPA₁ channels facilitate glycinergic transmission onto medullary dorsal horn neurons [25]. Considering that glutamate and glycine act as co-agonists of N-methyl-D-aspartate (NMDA) channels, this suggests that activation of TRPA₁ channels can lead to activation of NMDA channels. Although NMDA channels in the CeA account for maintenance of neuropathic pain [3], the contribution of TRPA₁ channels to pronociceptive changes in the CeA remains unknown.

From the canonical subfamily of TRP channels (TRPC), the amygdala expresses type 1, 4, and 5, which unite to form two heteromeric complexes: TRPC₁/TRPC₄ and TRPC₁/TRPC₅. In the dorsal root ganglion of the sciatic nerve, expression of TRPC₄ channels changes bidirectionally, depending on the site of injury and the outcome (pain or regeneration) [153,183]. In the LA, activation of mGLUR1 or mGLUR5 triggers an intracellular signaling cascade that leads to activation of TRPC₄ and TRPC₅ channels responsible for mediating innate but not learned (i.e., pain-related) fear [38,135,136]. In the CeA of neuropathic animals, however, it remains unknown whether TRPC₄ and TRPC₅ channels are responsible for mediating pain hypersensitivity and pain affect.

1.5 Somatosensory cortex

As the outer layer of the brain, the cortex provides an opportunity for external modulation of brain activity with minimum invasion. Since 1991, electrical stimulation of the primary motor cortex (M1) serves as a safer and more efficacious method for attenuation of chronic pain in patients than stimulation of the thalamus [163,164]. Nowadays, TMS of the M1 provides a minimally invasive alternative to direct electrical stimulation and serves as a method for treatment of two comorbid

deceases, chronic pain [92,110] and paresis [11,163]. However, TMS of the S2 can serve as an alternative and efficient method for attenuation of neuropathic pain, when TMS of the M1 remains inefficacious [99,100,168].

The human primary (S1) and secondary (S2) somatosensory cortices receive nociceptive and non-nociceptive somatosensory information directly from the thalamus and process it in parallel [98]. The S2 displays functional topography for noxious and innocuous stimuli [39]. Furthermore, innocuous stimulation of the nearby skin can inhibit (i.e., gate) nociceptive activation of the S2, whereas noxious stimulation cannot [161]. This is consistent with the proposal by Melzack and Wall [108] that innocuous stimulation may predetermine responses of the S1/S2 to noxious stimulation and trigger S1/S2-driven modulation of the gate control system in the spinal cord (see chapter 1.2.2).

In comparison with healthy controls, patients with trigeminal neuralgia [118] or central post-stroke pain [81] display a smaller volume of the gray matter in the S2, whereas patients with chronic musculoskeletal pain show greater neural activity in the S2 during presentation of pain-related words [159]. TMS of the S2 attenuates acute pain in healthy subjects [100,168] and drug-resistant neuropathic pain in patients [99]. Importantly, in both cases, the TMS of the S2 produces greater pain attenuation than TMS of the S1 or M1. In addition, TMS of the S2 attenuates visceral pain clinically [43], but does not, however, attenuate experimental pain in interictal migraineurs [165].

In healthy rats, electrical stimulation of the S2 fails to attenuate pain during acute noxious stimulation, whereas in rats with formalin-induced pain, electrical stimulation of the S2 produces weak antinociception by itself and strong antinociception in combination with a neural NO synthase inhibitor at a subtherapeutic dose [86]. The synergism of the two treatments correlates with c-Fos expression in the spinal cord and, in part, requires activation of spinal 5-HT receptors (but not activation of opioid and noradrenergic systems) [87]. Moreover, following formalin lip injections, S2 stimulation alone can reduce c-Fos expression in dorsal-horn neurons at the medullary level, suggesting its efficacy against orofacial pain [49]. Taken together, these findings indicate that S2 stimulation is capable of attenuating pain through medullospinal 5-HT mechanisms, but only in the setting of tissue injury.

2 AIMS

The aim of the current work was to test following hypotheses:

- Neuropathic pain results in part from pronociceptive hyperactivity of the CeA. Therefore, inhibition of the CeA with intraamygdaloid injections of lidocaine should attenuate neuropathic pain in rats with peripheral nerve injury but not nociceptive pain in sham-operated rats (**IV**). Furthermore, stimulation of the CeA with intraamygdaloid injections of glutamate should exacerbate neuropathic pain (**I**).
- Oxidative changes in the CeA contribute to development of pronociceptive hyperactivity of the CeA. Therefore, intraamygdaloid injections of antioxidants should attenuate neuropathic pain in rats with peripheral nerve injury, but not nociceptive pain in sham-operated rats (**IV**). Furthermore, intraamygdaloid injections of a ROS donor should recapitulate neuropathic pain characteristics (e.g. hypersensitivity) in sham-operated rats (**IV**).
- TRPC_{4/5} and TRPA₁ channels in the CeA act as receptors for ROS (for a review, see [121]). Therefore, intraamygdaloid injections of either a TRPC_{4/5} antagonist (**II**) or a TRPA₁ antagonist (**IV**) should attenuate neuropathic pain in rats with peripheral nerve injury, but not nociceptive pain in sham-operated rats.

In addition, we set to identify medullospinal mechanisms for hypothesized changes in spinal nociception induced by intraamygdaloid injections (**I**, **IV**). Particularly, we attempted to prevent hypothesized changes either by inhibiting cell bodies of 5-HT neurons in the RVM (**I**) or by blocking 5-HT receptors in the spinal cord (**I**, **IV**).

We next investigated whether S2 stimulation (**III**) attenuates neuropathic pain by reversing medullospinal mechanisms that the CeA engages for descending facilitation of spinal nociception (**I**, **IV**). We assessed the effect of S2 stimulation on neuropathic and nociceptive pain and recorded the discharge rate of medullary and spinal neurons in lightly anesthetized rats. Thereafter, we assessed whether the same drugs that prevented pain attenuation induced by intraamygdaloid injections (**I**, **IV**) would prevent pain attenuation induced by S2 stimulation (**III**).

3 METHODS

3.1 Ethical considerations

All experiments (Table 2) were performed according to the guidelines of European Communities Council Directive of 22 September 2010 (2010/63/EU). The national Animal Experiment Board in Finland (ELLA) approved the methods. All efforts were made to limit distress and to use only the number of rats (RccHan:WIST, Harlan Laboratories, Netherlands) necessary to produce reliable scientific data. We followed the principles of the 3Rs (replacement, reduction, and refinement) in the design of our experiments [146].

Table 2. Allocation of methods to their corresponding studies.

		Study
Surgeries		
	SNI model	I, II, IV
	SNL model	III
	Intracranial cannula implantation	I, II, III, IV
	Intrathecal catheterization	I, IV
Behavioral tests		
	Mechanical pain	I, II, III, IV
	Heat pain	III
	Emotional pain	IV
Electrophysiology		III

SNI spared nerve injury
 SNL spinal nerve ligation

3.2 Drugs

Glutamate (l-glutamic acid monosodium salt), 8-OH-DPAT (8-hydroxy-2-(dipropylamino)tetralin, 5-HT_{1A} receptor agonist), MK-801 (NMDA receptor antagonist), ML-204 (TRPC_{4/5} channel antagonist), naloxone (opioid receptor antagonist), ondansetron (5-HT₃ receptor antagonist), PBN (phenyl-N-tert-butyl-nitron, antioxidant), raclopride (dopamine D₂ receptor antagonist), t-BOOH (tert-butyl-hydroperoxide, ROS donor), TEMPOL (4-hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl, antioxidant), WAY-100635 (5-HT_{1A} receptor antagonist), and A-967079 (TRPA₁ channel antagonist) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Atipamezole (α_2 -adrenoceptor antagonist), lidocaine (sodium channel blocker), and pentobarbital (GABA_A receptor positive allosteric modulator) were purchased from OrionPharma (Espoo, Finland). CHEMBRIDGE-5861528 (TRPA₁ channel antagonist) was purchased from ChemBridge (San Diego, CA, USA). Buprenorphine (μ -receptor partial agonist, κ -receptor antagonist) was purchased from RB Pharmaceuticals Limited (Slough, UK).

3.3 Surgeries

3.3.1 Anesthesia

All surgeries were performed in rats anesthetized with intraperitoneal injections of pentobarbital at the dose of 60 mg per kg. Additional injections of pentobarbital were given at the dose of 15-20 mg per kg as needed to maintain deep level of anesthesia. Following surgery, rats received subcutaneous injections of buprenorphine at the dose of 0.01-0.03 mg per kg for three consecutive days every twelve hours. Rats were allowed to recover for at least one week before the experiments (Fig. 4).




Figure 4. Time course of studies I-IV.

3.3.2 Spared nerve injury model

The spared nerve injury (SNI; Fig. 5) model of neuropathic pain derives its name from sparing the sural nerve while cutting the other two branches of the sciatic nerve (tibial and common peroneal nerves) [30]. Such partial denervation induces pain hypersensitivity in the lateral plantar surface of the injured paw for six or more months.




Figure 5. Spared nerve injury model.

3.3.3 Spinal nerve ligation model

Unilateral ligation of spinal nerves L5 and L6 served as a second model of neuropathic pain [78]. In contrast to SNI, rats with spinal nerve ligation (SNL; Fig. 6) develop pain hypersensitivity in the entire plantar surface of the injured paw. Hypersensitivity in the SNL model, however, lasts for a much shorter duration (up to two months).




Figure 6. Spinal nerve ligation model.

3.3.4 Intracranial cannula implantation

Anesthetized rats were secured to a stereotaxic frame and a hole was made in the skull for the implantation of a chronic guide cannula 2 mm above the desired injection site. The guide cannula was cemented to screws, which, in turn, were anchored to the skull. During experiments, injections were made with the use of an internal cannula protruding 2 mm from the tip of the guide cannula. At the completion of experimental procedures, the rats were euthanized, and injection sites were identified in the coronally sectioned brain after fixation in formalin.

3.3.5 Intrathecal catheterization

Anesthetized rats were implanted with an intrathecal catheter for lumbar drug injections. The implantation was considered successful if a lidocaine injection through the catheter produced temporal paralysis of hind limbs.

3.4 Behavioral tests

3.4.1 Mechanical pain

Rate of paw withdrawal during mechanical stimulation with ascending series of calibrated von Frey filaments indicated the level of mechanical pain sensitivity.

3.4.2 Heat pain

Latency of paw withdrawal during heat stimulation with a Peltier thermode indicated the level of heat pain sensitivity.

3.4.3 Emotional pain

To evaluate the aversive component of pain, an aversive place-conditioning paradigm was used [89], in which a neuropathic rat resided in a chamber divided into two halves, one dark and the other illuminated. In the dark half, the injured paw was stimulated, whereas, in the illuminated half, the healthy paw was stimulated. As a result, the rat chose the illuminated half over the more naturally preferable dark half. A drug-induced increase in time the rat spent in the dark half indicated attenuation of emotional pain.

3.5 Electrophysiology

The discharge rate of neurons was recorded with an extracellular electrode in lightly anesthetized rats secured to a stereotaxic frame. In the RVM, neurons giving excitatory responses to noxious stimulation were considered as pronociceptive ON cells, whereas those giving inhibitory responses were considered as antinociceptive OFF cells [56]. In the SDH, neurons giving excitatory responses to both noxious and innocuous stimulation were considered as pain-relaying WDR cells.

4 RESULTS AND DISCUSSION

4.1 Medullospinal 5-HT neurons as a relay for bidirectional modulation of pain hypersensitivity by the amygdala (I, IV)

As a first step in investigating the role of the amygdala in neuropathic pain, we studied how inhibition of the right CeA with lidocaine influences pain in neuropathic and non-neuropathic conditions. We chose the right CeA as it plays a greater role in the development of inflammatory pain than the left CeA, independent of the location of inflammation [23,70]. Furthermore, the right CeA was contralateral to the nerve injury. A previous study in similar settings demonstrates greater pronociceptive changes in the right CeA than in the left one [65]. In our experiments, lidocaine injection (4 %, 0.5 µl) into the CeA attenuated pain hypersensitivity in neuropathic rats (SNI model) but not nociceptive pain in sham-operated rats (Table 3). This indicates that pain hypersensitivity in peripheral neuropathy develops, at least in part, due to pronociceptive hyperactivity of the CeA. This is consistent with other studies, which demonstrate positive correlation of amygdaloid activity with the level of chronic pain [44,47,65,71,85,116].

However, because lidocaine inhibits both neurons that originate from the CeA and neurons that pass through the CeA, our second step was to investigate how pain hypersensitivity changes during modulation of synaptic activity in the CeA with various doses of glutamate. A high dose of glutamate (100 µg) in the CeA attenuated hypersensitivity, whereas a low dose of glutamate (9 µg) facilitated it. Furthermore, blocking NMDA receptors in the CeA with MK-801 prevented facilitation of hypersensitivity induced by a low dose of glutamate but not attenuation of hypersensitivity induced by a high dose of glutamate. In these experiments, we pretreated the CeA with MK-801 at a dose that was too low to influence pain hypersensitivity by itself. However, in another set of experiments, attenuation of pain hypersensitivity was evident after the dose of MK-801 was elevated. Taken together, this suggests that, in the CeA of neuropathic animals, endogenous release of glutamate causes activation of NMDA receptors that dominates over activation of non-NMDA glutamate receptors. As a result, the CeA displays pronociceptive hyperactivity in neuropathy. This is consistent with a previous study that demonstrates contribution of NMDA receptors in the CeA to development of pain hypersensitivity [3]. On the other hand, there is also evidence that non-NMDA glutamate receptors in the CeA modulate pain hypersensitivity bidirectionally, depending on their type and site of expression [54,96,116,134]. However, it was outside the scope of this thesis to identify which non-NMDA

glutamate receptors in the CeA mediated attenuation of hypersensitivity induced by a high dose of glutamate.

To determine if CeA treatment with a high dose of glutamate attenuates hypersensitivity by exerting its action at the spinal cord level, we attempted to prevent antinociceptive action of glutamate by blocking spinal receptors, which are normally associated with antinociception (for a review, see [111]). Blocking 5-HT_{1A} receptors with WAY-100635 prevented attenuation of hypersensitivity induced by a high dose of glutamate in the CeA, whereas blocking α_2 -adrenoceptors (with atipamezole), or dopamine D₂ receptors (with raclopride), or opioid receptors (with naloxone) failed to do the same. Furthermore, spinal injections by themselves had no effect on hypersensitivity at the currently used doses. Considering that 5-HT in the spinal cord originates entirely from supraspinal regions [88], this indicates that the antihypersensitivity effect induced by a high dose of glutamate in the CeA is mediated primarily by spinally projecting 5-HT neurons. This is consistent with other studies, which demonstrate the importance of spinal 5-HT_{1A} receptors in mediating attenuation of hypersensitivity in neuropathic [179] and other pathological pain states [48,186].

To determine if spinally projecting 5-HT neurons also mediate facilitation of hypersensitivity induced by a low dose of glutamate in the CeA, we attempted to prevent the pronociceptive action of glutamate by blocking the spinal 5-HT₃ receptor, which is normally associated with pronociception (for a review, see [111]). Blocking 5-HT₃ receptors with ondansetron, which alone had no effect on hypersensitivity (i.e., at the currently used dose), prevented facilitation. This indicates that spinally projecting 5-HT neurons in the medulla can relay both facilitation and attenuation of hypersensitivity from the CeA. Although this is consistent with other studies, which demonstrate contribution of spinal 5-HT₃ receptors to the development of hypersensitivity, there is also evidence of their role in mediating antihypersensitivity. Such controversy may result from the fact that activation of spinal 5-HT₃ receptors can lead to the release of spinal GABA [55], the inhibitory potency of which can disappear at certain stages of chronic pain development [26,27].

We next investigated whether medullospinal neurons act as sources of 5-HT in the spinal cord for bidirectional modulation of hypersensitivity by the CeA. We were particularly interested in the role of the RVM, which accounts for antinociception in pain-related fear mediated by the amygdala [58,59]. We therefore pretreated the RVM with a 5-HT_{1A} autoreceptor agonist (8-OH-DPAT) in an attempt to hyperpolarize cell bodies of 5-HT neurons [142]. Although RVM pretreatment with 8-OH-DPAT by itself failed to influence hypersensitivity, 8-OH-DPAT in the RVM prevented both facilitation of hypersensitivity induced by a low dose of glutamate

and attenuation of hypersensitivity induced by a high dose of glutamate in the CeA. This indicates that the CeA regulates activation of spinal 5-HT receptors through a release of 5-HT from neurons, which descend to the spinal cord from the RVM. However, it remains unknown how the CeA exerts its bidirectional action on hypersensitivity by activating the same medullospinal 5-HT neurons. It is possible that the CeA selectively activates either pronociceptive 5-HT₃ receptors or antinociceptive 5-HT_{1A} receptors by activating different subtypes of medullospinal 5-HT neurons. Alternatively, the CeA may regulate the level of net activity of medullospinal 5-HT neurons and, thereby, titrate the release of 5-HT in the spinal cord. This notion is supported by the findings that treatment of the RVM with a low dose of glutamate leads to facilitation of nociceptive pain, whereas treatment with a high dose results in attenuation [187]. Moreover, 5-HT neurons in the RVM account for both weak and strong attenuations of nociceptive pain induced by treatment of the PAG with low and high doses of morphine, respectively [17].

In conclusion, endogenous release of glutamate predominately leads to activation of NMDA receptors, which maintain pronociceptive hyperactivity of the CeA. The CeA, in turn, exerts its pronociceptive hyperactivity by preventing antinociceptive activation of medullospinal 5-HT neurons.

Table 3. Medullospinal 5-HT neurons as a relay for bidirectional modulation of pain hypersensitivity by the amygdala.

		CeA injections		
		LIDO (IV)	GLU100 (I)	GLU9 (I)
Neuropathic rats (SNI)	Action in the CeA	block of sodium channels	activation of non-NMDA receptors	activation of NMDA receptors
	Action in the RVM	–	activation of 5-HT cell bodies	activation of 5-HT cell bodies
	Action in the SDH	–	activation of 5-HT _{1A} receptors	activation of 5-HT ₃ receptors
	Pain hypersensitivity	↓	↓	↑
	Pain affect	–	–	–
Sham rats	Nociceptive pain	0	–	–

CeA	central amygdaloid nucleus	↑	exacerbation
GLU100	100 µg of glutamate	↓	attenuation
GLU9	9 µg of glutamate	0	no effect
LIDO	lidocaine	–	not studied
RVM	rostral ventromedial medulla		
SDH	spinal dorsal horn		
SNI	spared nerve injury model		

4.2 Reactive oxygen species as a culprit for pronociceptive hyperactivity of the amygdala (II, IV)

In light of recent findings [97] that glutamate in the CeA can exacerbate nociceptive pain by promoting production of ROS, we addressed the role of ROS in the development of neuropathic pain. To test this, we first assessed how treatment of the CeA with antioxidants (TEMPOL and PBN) influences pain in neuropathic and non-neuropathic conditions. Treatment of the CeA with either of the two antioxidants attenuated pain hypersensitivity in neuropathic rats (SNI model). In sham-operated rats, however, neither TEMPOL nor PBN influenced nociceptive pain (Table 4). Thereafter, we addressed whether attenuation of pain affect accompanies attenuation of hypersensitivity in neuropathic rats after treatment of the CeA with antioxidants. We chose TEMPOL from the two antioxidants and observed that it attenuates pain affect. This suggests that neuropathy induces oxidative changes in the CeA that may account for its pronociceptive and pain-affect-inducing hyperactivity. This is consistent with other studies demonstrating contribution of ROS in the CeA to the development of visceral pain and bee venom-induced pain [68,101].

Because ROS act as endogenous agonists of TRP channels (for a review, see [121]), we next investigated if TRPC_{4/5} and TRPA₁ channels in the CeA mediate observed oxidative changes. We therefore assessed how treatment of the CeA with antagonists of a TRPC_{4/5} channel (ML-204) and of a TRPA₁ channel (CHEMBRIDGE-5861528) influences neuropathic pain. Similar to antioxidants, ML-204 and CHEMBRIDGE-5861528 in the CeA separately attenuated both pain hypersensitivity and pain affect in neuropathic rats but not nociceptive pain in sham-operated rats. In addition, attenuation of hypersensitivity was evident after blocking TRPA₁ channels with an antagonist structurally different from CHEMBRIDGE-5861528, A-967079. Taken together, this indicates that, in neuropathy, ROS account for pronociceptive and pain affect-inducing hyperactivity of the CeA by acting on TRPC_{4/5} and TRPA₁ channels. This is somewhat contrary to other studies, in which TRPC_{4/5} channels in the amygdala failed to play a role in pain-related fear [135,136]. However, this controversy can be explained by the fact that these studies investigated the role of TRPC_{4/5} channels in the LA, whereas we focused on their role in the CeA. Additionally, these studies [135,136] demonstrate inability of TRPC_{4/5} channels to influence pain-related fear in healthy (knockout) animals, whereas we examined the ability of TRPC_{4/5} channels to influence pain affect only in neuropathic animals. Considering, however, that, in our experiments, blocking TRPC_{4/5} channels failed to attenuate nociceptive pain in non-neuropathic (sham-

operated) animals, it is likely that TRPC_{4/5} channels account for pain affect and pain-related fear only in the setting of nerve injury.

Since activation of 5-HT_{1A} receptors in the spinal cord accounted for attenuation of hypersensitivity induced by a high dose of glutamate in the CeA, we next investigated whether they also account for attenuation of hypersensitivity seen after blocking TRPA₁ channels in the CeA with CHEMBRIDGE-5861528. As expected, blocking spinal 5-HT_{1A} receptors with WAY-100635 at a dose that alone failed to influence hypersensitivity prevented the antihypersensitivity effect of CHEMBRIDGE-5861528. Considering that a high dose of glutamate in the CeA attenuates hypersensitivity by activating medullospinal 5-HT neurons, this suggests that blocking TRPA₁ channels in the CeA leads to a similar activation of medullospinal 5-HT neurons.

Encouraged by above findings, we attempted to recapitulate hypersensitivity in sham-operated rats by treating the CeA with a ROS donor, t-BOOH. Although, t-BOOH in a previous study exacerbated nociceptive pain after administration into the spinal cord [180], it failed to do so in the current study after administration at the same dose into the CeA. This indicates that acute administration of ROS alone is insufficient to induce pronociceptive changes in the CeA and thus suggests that its chronic administration or presence of comorbid factors (or both) is necessary.

In conclusion, ROS prevent antinociceptive activation of medullospinal 5-HT neurons by promoting pronociceptive hyperactivity of the CeA.

Table 4. Reactive oxygen species as a culprit for pronociceptive hyperactivity of the amygdala.

		CeA injections				
		TEMPOL (IV)	PBN (IV)	ML-204 (II)	CHEM (IV)	t-BOOH (IV)
Neuropathic rats (SNI)	Action in the CeA	anti- oxidant	anti- oxidant	block of TRPC _{4/5} channels	block of TRPA ₁ channels	ROS donor
	Action in the RVM	–	–	–	–	–
	Action in the SDH	–	–	–	activation of 5-HT _{1A} receptors	–
	Pain hyper- sensitivity	↓	↓	↓	↓	–
	Pain affect	↓	–	↓	↓	–
Sham rats	Nociceptive pain	0	0	0	0	0

CeA	central amygdaloid nucleus	↓	attenuation
CHEM	CHEMBRIDGE-5861528	0	no effect
PBN	phenyl-N-tert-butyl-nitrone	–	not studied
ROS	reactive oxygen species		
RVM	rostral ventromedial medulla		
SDH	spinal dorsal horn		
SNI	spared nerve injury model		
t-BOOH	tert-butyl-hydroperoxide		
TEMPOL	4-hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl		

4.3 Medullospinal 5-HT neurons as a relay for attenuation of neuropathic pain by cortical stimulation (III)

TMS of the S2 has proven to be an efficacious method for attenuation of drug-resistant pain in neuropathic patients [99]. However, because the mechanism of pain attenuation remains largely unknown, we examined whether descending pathways, which are activated by S2 stimulation, overlap with those pathways that are activated by a high dose of glutamate in the CeA. We therefore investigated whether medullary 5-HT or non-5-HT neurons contribute to S2 stimulation-induced pain attenuation in neuropathic rats. To test this, we recorded neurons in the RVM in parallel with behavioral assessment of pain. In addition, we separately recorded pain-relay neurons in the SDH. However, because SDH recordings required irrecoverable surgical preparations (laminectomy), we performed them in lightly anesthetized rats. Furthermore, to obtain comparable data, assessment of pain behavior and simultaneous recordings of medullospinal neurons were also performed in lightly anesthetized rats. However, because light anesthesia diminishes the withdrawal reflex, we used a piezoelectric movement detector over the gastrocnemius muscle to assess the reflex mechanomyographically. Furthermore, we replaced noxious mechanical stimulation with noxious heat stimulation to eliminate artifacts in the mechanomyogram. This, in turn, required implementation of the SNL model of neuropathic pain instead of the SNI model to enlarge the hypersensitive surface of the paw at the expense of the duration of hypersensitivity: two months in SNL rats versus six (or more) months in SNI rats [30,78].

Stimulation of the S2, independent of the hemisphere, attenuated heat pain in nerve-injured rats with mechanical hypersensitivity. Attenuation of heat pain was evident at the lowest cortical stimulus intensity of 30 μ A (0.1-ms pulses at 300 Hz). The antinociceptive effect failed to increase at higher intensities of 50 and 70 μ A. This suggests that pain attenuation was induced by stimulation of the S2 and not by a possible spread of electrical current to adjacent regions. Stimulation of the right S2 (contralateral to the injury), however, failed to attenuate heat pain in sham-operated rats and in rats that failed to develop mechanical hypersensitivity after SNL (Table 5). Although it remains unknown whether S2 stimulation in the left hemisphere attenuates pain in sham-operated and non-hypersensitive neuropathic rats, this suggests that S2 stimulation, at least in the right hemisphere, is efficacious only in animals with pathological pain hypersensitivity. This is consistent with previous findings that S2 stimulation attenuates inflammatory but not nociceptive pain [64,86].

Thereafter, we performed recordings in the SDH during stimulation of the right S2 to investigate spinal mechanisms for behavioral attenuation of heat pain in hypersensitive neuropathic rats. In particular, we recorded wide-dynamic range (WDR) cells, which accelerated their discharge rate in response to both noxious heat stimulation and innocuous mechanical stimulation of the injured paw. We observed that S2 stimulation attenuates heat responses of WDR cells (Fig. 7). We observed also that this attenuation can be prevented by blocking spinal 5-HT_{1A} receptors with WAY-100635 at a dose that alone failed to reduce heat-response discharge. Considering that, at the current dose, WAY-100635 also failed to reduce the ongoing discharge rate of WDR cells, this suggests that S2 stimulation attenuated pain by activating 5-HT_{1A} receptors, which are presynaptic to WDR cells. However, one must be cautious with this interpretation, since failure of WAY-100635 to reduce the ongoing discharge rate of WDR cells may have resulted from the fact that the ongoing discharge rate of WDR cells was originally very low. On the other hand, a recent whole-cell patch clamp study in spinal cord slices supports our interpretation of presynaptic localization of 5-HT_{1A} receptors, by demonstrating that 5-HT in the SDH inhibits nociceptive transmission via activation of 5-HT_{1A} receptors, which are localized on spinal terminals of primary afferents [160].




Figure 7. Example of recordings in the spinal dorsal horn.

Because both electrical stimulation of the S2 and glutamatergic stimulation of the CeA attenuate pain in hypersensitive neuropathic rats by activating spinal 5-HT_{1A} receptors, it is likely that they do so through the same mechanism. To test this hypothesis, we attempted to prevent S2 stimulation-induced pain attenuation by pretreating the RVM with an agonist of 5-HT_{1A} autoreceptors (8-OH-DPAT), which prevented pain attenuation induced by a high dose of glutamate in the CeA. In the S2 experiments, as in the CeA experiments, we used 8-OH-DPAT at a dose, which failed to influence pain by itself. In support of our hypothesis, 8-OH-DPAT prevented S2 stimulation-induced pain attenuation. As in the CeA experiments, we presume that 8-OH-DPAT acted by inhibiting cell bodies of spinally projecting 5-HT neurons. This indicates that activation of medullospinal 5-HT neurons is a necessary step for the S2 stimulation-induced pain attenuation, as it is for pain attenuation induced by a high dose of glutamate in the CeA. However, it remains unknown whether S2 stimulation and a high dose of glutamate in the CeA attenuate pain by recruiting the same subtypes of medullospinal 5-HT neurons or by regulating their net activity in a similar fashion.




Figure 8. Example of recordings in the rostral ventromedial medulla and parallel assessment of pain behavior.

In addition, we recorded heat responses of pain-regulating ON and OFF cells in the RVM, which are shown to be non-5-HT [106]. Unlike WDR cells in the SDH, which respond to both noxious and innocuous stimulations, ON and OFF cells in the RVM respond only to noxious stimulation. Furthermore, ON cells respond by accelerating their discharge rate before withdrawal of the paw, whereas OFF cells respond by decelerating it. A fast discharge rate of ON cells and a slow discharge rate of OFF cells predict facilitation of nociception, whereas a slow discharge rate of ON cells and a fast discharge rate of OFF cells predict inhibition of nociception [56]. However, in our experimental setup, we were unable to record ON and OFF cells during S2 stimulation, because the cortical stimulation electrode induced stimulus artifacts, which masked all neural spikes during stimulation. Therefore, we were able to record ON and OFF cells only after cessation of S2 stimulation when attenuation of pain behavior was marginal. Nonetheless, we observed that S2 stimulation shortens duration of heat responses in ON cells and prolongs the latency of heat responses in OFF cells (Fig. 8). Considering that at least some of the ON and OFF cells project to the SDH [40,169], this suggests that S2 stimulation promotes antinociceptive activity of both non-5-HT (i.e., ON and OFF cells) and 5-HT medullospinal neurons. Therefore, our findings are consistent with a previous study [87], where unselective inhibition of spinal 5-HT receptors prevented pain attenuation induced by S2 stimulation. Interestingly, the same study [87] demonstrates also the insignificance of the opioid and noradrenergic systems in S2 stimulation-induced pain attenuation, which narrows the list of candidates for the role of (presumably medullospinal) non-5-HT neurons engaged by S2 stimulation in our study.

In conclusion, S2 stimulation represents a clinically relevant method that has the potential to recover antinociceptive activity of medullospinal 5-HT neurons, which is suppressed in neuropathy due to pronociceptive hyperactivity of the CeA.

Table 5. Medullospinal 5-HT neurons as a relay for attenuation of neuropathic pain by cortical stimulation.

		S2 stimulation (III)
Neuropathic rats (SNL)	Action in the CeA	–
	Action in the RVM	activation of 5-HT cell bodies
		inhibition of ON cell
		activation of OFF cells
	Action in the SDH	inhibition of WDR cells via activation of 5-HT _{1A} receptors
	Pain hyper-sensitivity	↓
	Pain affect	–
Sham rats	Nociceptive pain	0

CeA	central amygdaloid nucleus	↓	attenuation
RVM	rostral ventromedial medulla	0	no effect
SDH	spinal dorsal horn	–	not studied
SNL	spinal nerve ligation model		
S2	secondary somatosensory cortex		
WDR	wide-dynamic range		

4.4 General discussion

The present studies suggest that, in neuropathic conditions, endogenous release of glutamate in the CeA predominantly activates NMDA (rather than non-NMDA) receptors, which maintain pronociceptive hyperactivity of the CeA (I). However, activation of non-NMDA glutamate receptors in the CeA can lead to attenuation of pain hypersensitivity, which is likely due to suppression of pronociceptive hyperactivity of the CeA. CeA-induced pronociception results from activation of spinal 5-HT₃ receptors, whereas CeA-induced antinociception results from activation of spinal 5-HT_{1A} receptors. Interestingly, activation of both spinal 5-HT₃ and 5-HT_{1A} receptors requires activation of medullospinal 5-HT neurons. This indicates that the CeA normally promotes antinociceptive activity of medullospinal 5-HT neurons, whereas, in neuropathy, the CeA starts to promote their pronociceptive activity. It is likely that medullospinal 5-HT neurons have two distinct subtypes, one of which relays antinociception, whereas the other accounts for pronociception, similar to medullospinal non-5-HT neurons (antinociceptive OFF cells and pronociceptive ON cells) [40,56,106,169]. Alternatively or additionally, the level of net activity of medullospinal 5-HT neurons may determine the final effect on nociception.

We also provide evidence that ROS in the CeA can promote pronociceptive hyperactivity of the CeA through activation of TRPA₁ and TRPC_{4/5} channels (II, IV). In light of these observations, an increase in ROS production and a decrease in ROS detoxification in the CeA emerge as possible maladaptive changes that account for the development of neuropathic pain. Furthermore, considering that ROS represent a product of mitochondrial activity (for a review, see [113]), neuropathic pain can in part represent a mitochondrial disorder. This is consistent with previous findings that chemotherapy induces neuropathic pain by promoting accumulation of atypical mitochondria in peripheral nerves, which is associated with ROS overproduction and poor ROS detoxification [8,184].

In addition, we established that S2 stimulation attenuates neuropathic pain by inhibiting spinal transmission of nociceptive information through WDR cells (III). To accomplish this, S2 stimulation employs medullospinal 5-HT neurons to activate spinal 5-HT_{1A} receptors, which are likely presynaptic to WDR cells. Interestingly, such mechanism for pain attenuation in SNL rats is similar to the one that the CeA engages in SNI rats. This indicates that malfunction of medullospinal 5-HT neurons may be an important contributor to the development of chronic pain in peripheral neuropathies. Furthermore, this suggests that S2 stimulation acts by reversing CeA-driven pronociceptive activation of medullospinal 5-HT neurons. Other mechanisms for S2 stimulation-induced pain attenuation include antinociceptive changes in

activity of (presumably medullospinal) non-5-HT neurons (ON and OFF cells) [40,56,106,169].

5 CAVEATS

Although intracranial cannulae targeted the CeA (**I**, **II**, **IV**), the volume of injections was large enough (0.5 μ l) for drugs to spread to adjacent brain structures, including other amygdaloid nuclei. Therefore, the action of drugs on neurons outside the CeA may have interfered with our results. On the other hand, the CeA acts as the output nucleus of the amygdala with regard to pain modulation. Furthermore, injection of drugs into a control site (internal capsule) failed to produce any alterations in pain behavior. Taken together, this indicates that changes in pain behavior, most likely, resulted from the action of drugs on neurons in the CeA.

Caution is necessary in comparing the results collected from the different studies in this thesis. In amygdaloid studies (**I**, **II**, **IV**), we assessed mechanical pain in freely moving SNI rats. In the cortical study (**III**), we assessed heat pain in lightly anesthetized SNL rats. Such inconsistency in experimental settings may have introduced bias to our results, which demonstrate overlap in mechanisms for pain regulation by the amygdala (i.e., CeA) and the cortex (i.e., S2). Furthermore, not all aspects of pain were assessed. In studies **II** and **IV**, we assessed sensory-discriminative and affective-motivational aspects of pain, whereas in studies **I** and **III**, we assessed only the former. We also did not assess ongoing pain in any of the studies, although ongoing pain represents a major complaint among neuropathic patients.

We inhibited medullospinal 5-HT neurons by activating 5-HT_{1A} autoreceptors on their cell bodies with 8-OH-DPAT injections into the RVM (**I**, **III**). We, however, cannot exclude that activation of 5-HT_{1A} heteroreceptors on other cells (neural [142] or glial [5]) influenced our results.

Although previous studies demonstrate that at least some of the ON and OFF cells in the RVM have spinal projections [40,169], we did not verify that the ON and OFF cells recorded in our study (**III**) indeed represented medullospinal neurons. Furthermore, the RVM can comprise ascending projections from the area around the central canal of the spinal cord [62]. If these ascending projections contribute to transmission of nociception, then they may resemble electrophysiologically ON or OFF cells. Collision testing, for example, could have clarified whether recorded in our study (**III**) RVM cells were ascending or descending.

6 FUTURE PROSPECTS

We demonstrated that pain-modulating pathways descending from the S2 (III) and the CeA (I, IV) merge in the RVM. It is therefore possible that the S2 and the CeA modulate pain in tandem. Future studies may address whether S2 stimulation attenuates neuropathic pain by modulating the activity of the CeA or by modulating the activity of the PAG, which connects the CeA to the RVM.

We also demonstrated that administration of antioxidants or TRPA₁ antagonists into the CeA attenuates neuropathic pain (IV), which is consistent with previous findings of their antinociceptive properties in the spinal cord [178,180]. From a clinical perspective, it is therefore of great interest to see if antioxidants and TRPA₁ antagonists can attenuate neuropathic pain, when administered systemically, and to examine whether their action at the spinal or supraspinal level plays a more important role in attenuation of neuropathic pain. In addition, it is interesting to see if systemic administration of antioxidants can potentiate antinociceptive action of S2 stimulation.

A recent clinical study [99] demonstrated that the S2 is an effective cortical stimulation site for attenuating neuropathic pain with TMS. In line with this, our study (III) shows that electrical stimulation of the S2 also attenuates neuropathic pain in rats and reveals the mechanism for S2 stimulation-induced pain attenuation. It remains to be studied whether a similar mechanism contribute to pain attenuation induced by TMS of the S2 in neuropathic patients.

Implementation of antidepressants, such as selective serotonin reuptake inhibitors (SSRI), has produced controversial results in treatment of neuropathic pain (for a review, see [82]). This is consistent with our findings, which suggest that medullospinal 5-HT neurons can account for both attenuation (I, III, IV) and exacerbation (I) of neuropathic pain. Investigation of how various SSRI influence the activity of medullospinal 5-HT neurons may lead to improved clinical use.

Since electrical stimulation of the S2 (III) and pharmacological inhibition of the CeA (I, IV) attenuated neuropathic pain by activating spinal 5-HT_{1A} receptors, it may be expected that spinal administration of a 5-HT_{1A} agonist also attenuates neuropathic pain. However, since pain attenuation induced by electrical stimulation of the S2 and by pharmacological inhibition of the CeA can be prevented by activation of 5-HT_{1A} receptors in the RVM (I, III), systemic administration of a 5-HT_{1A} agonist may prove inefficacious. In light of this notion, spinal (but not systemic) administration of BUSPIRONE may serve as a new method for clinical treatment of neuropathic pain. BUSPIRONE represents one of the few clinically approved drugs that acts as a 5-HT_{1A} receptor agonist. Currently, BUSPIRONE is used to treat anxiety but not

neuropathic pain, despite the fact that anxiety represents one of the comorbidities of neuropathic pain. The reason for this lies in inefficacy of BUSPIRONE to attenuate pain in neuropathic patients when administered systemically [79]. Our studies (**I**, **III**, **IV**), however, raise the hypothesis that spinal administration of BUSPIRONE may prove efficacious.

7 CONCLUSIONS

In neuropathic conditions:

- NMDA receptors in the CeA maintain its pronociceptive hyperactivity characterized by activation of spinal 5-HT₃ receptors (I).
- Activation of non-NMDA glutamate receptors in the CeA suppresses its pronociceptive hyperactivity and leads to descending antinociception mediated by spinal 5-HT_{1A} receptors (I).
- ROS contribute to CeA-driven pain hypersensitivity and pain affect through activation of TRPA₁ and TRPC_{4/5} channels in the CeA (II, IV).
- Activation of medullospinal 5-HT neurons is necessary for both CeA-induced facilitation and inhibition of spinal nociception (I, IV).
- S2 stimulation attenuates neuropathic pain by promoting antinociceptive activity of medullospinal 5-HT neurons and of (presumably medullospinal) non-5-HT neurons (ON and OFF cells; III).

By summarizing the results of the present studies (I-IV), the table below (Table 6) describes the mechanisms of neuropathic pain development and suggests new approaches to its treatment.

Table 6. Mechanisms of neuropathic pain development and possible approaches to its treatment.

		RVM	SDH	Pain
CeA	activation of NMDA receptors	pronociceptive activation of medullospinal 5-HT neurons	activation of 5-HT ₃ receptors	↑
	activation of TRPA ₁ & TRPC _{4/5} receptors with ROS			
CeA	block of NMDA receptors with antagonists	antinociceptive activation of medullospinal 5-HT neurons	activation of 5-HT _{1A} receptors	↓
	activation of non-NMDA glutamate receptors			
	block of TRPA ₁ & TRPC _{4/5} receptors with antagonists			
	detoxification of ROS with antioxidants			
S2	electrical stimulation	antinociceptive activation of medullospinal 5-HT neurons	activation of 5-HT _{1A} receptors	↓
			inhibition of WDR cells	
		antinociceptive activation of (likely medullospinal) non-5-HT neurons (ON and OFF cells)	–	

CeA	central amygdaloid nucleus	↑	exacerbation
ROS	reactive oxygen species	↓	attenuation
RVM	rostral ventromedial medulla	–	not studied
SDH	spinal dorsal horn		
S2	secondary somatosensory cortex		
WDR	wide-dynamic range		

8 REFERENCES

- [1] Agnati LF, Fuxe K, Zoli M, Ozini I, Toffano G, Ferraguti F. A correlation analysis of the regional distribution of central enkephalin and beta-endorphin immunoreactive terminals and of opiate receptors in adult and old male rats. Evidence for the existence of two main types of communication in the central nervous system: the volume transmission and the wiring transmission. *Acta Physiol Scand* 1986;128:201-207.
- [2] Andersson DA, Gentry C, Moss S, Bevan S. Transient receptor potential A1 is a sensory receptor for multiple products of oxidative stress. *J Neurosci* 2008;28:2485-2494.
- [3] Ansah OB, Bourbia N, Goncalves L, Almeida A, Pertovaara A. Influence of amygdaloid glutamatergic receptors on sensory and emotional pain-related behavior in the neuropathic rat. *Behav Brain Res* 2010;209:174-178.
- [4] Ansah OB, Leite-Almeida H, Wei H, Pertovaara A. Striatal dopamine D2 receptors attenuate neuropathic hypersensitivity in the rat. *Exp Neurol* 2007;205:536-546.
- [5] Azmitia EC, Gannon PJ, Kheck NM, Whitaker-Azmitia PM. Cellular localization of the 5-HT1A receptor in primate brain neurons and glial cells. *Neuropsychopharmacology* 1996;14:35-46.
- [6] Bakshi VP, Smith-Roe S, Newman SM, Grigoriadis DE, Kalin NH. Reduction of stress-induced behavior by antagonism of corticotropin-releasing hormone 2 (CRH2) receptors in lateral septum or CRH1 receptors in amygdala. *J Neurosci* 2002;22:2926-2935.
- [7] Bao L, Wang HF, Cai HJ, Tong YG, Jin SX, Lu YJ, Grant G, Hokfelt T, Zhang X. Peripheral axotomy induces only very limited sprouting of coarse myelinated afferents into inner lamina II of rat spinal cord. *Eur J Neurosci* 2002;16:175-185.
- [8] Barriere DA, Rieusset J, Chanteranne D, Busserolles J, Chauvin MA, Chapuis L, Salles J, Dubray C, Morio B. Paclitaxel therapy potentiates cold hyperalgesia in streptozotocin-induced diabetic rats through enhanced mitochondrial reactive oxygen species production and TRPA1 sensitization. *Pain* 2012;153:553-561.
- [9] Basso MA, Evinger C. An explanation for reflex blink hyperexcitability in Parkinson's disease. II. Nucleus raphe magnus. *J Neurosci* 1996;16:7318-7330.
- [10] Basso MA, Powers AS, Evinger C. An explanation for reflex blink hyperexcitability in Parkinson's disease. I. Superior colliculus. *J Neurosci* 1996;16:7308-7317.

- [11] Belci M, Catley M, Husain M, Frankel HL, Davey NJ. Magnetic brain stimulation can improve clinical outcome in incomplete spinal cord injured patients. *Spinal Cord* 2004;42:417-419.
- [12] Bernard JF, Besson JM. The spino(trigemino)pontoamygdaloid pathway: electrophysiological evidence for an involvement in pain processes. *J Neurophysiol* 1990;63:473-490.
- [13] Bhatnagar S, Vining C, Denski K. Regulation of chronic stress-induced changes in hypothalamic-pituitary-adrenal activity by the basolateral amygdala. *Ann N Y Acad Sci* 2004;1032:315-319.
- [14] Bird GC, Lash LL, Han JS, Zou X, Willis WD, Neugebauer V. Protein kinase A-dependent enhanced NMDA receptor function in pain-related synaptic plasticity in rat amygdala neurones. *J Physiol* 2005;564:907-921.
- [15] Black AK, Fulwiler JC, Smitherman TA. The role of fear of pain in headache. *Headache* 2015;55:669-679.
- [16] Blackburn-Munro G, Blackburn-Munro RE. Chronic pain, chronic stress and depression: coincidence or consequence? *J Neuroendocrinol* 2001;13:1009-1023.
- [17] Borszcz GS. Differential contributions of medullary, thalamic, and amygdaloid serotonin to the antinociceptive action of morphine administered into the periaqueductal gray: a model of morphine analgesia. *Behav Neurosci* 1999;113:612-631.
- [18] Bos R, Sadlaoud K, Boulenguez P, Buttigieg D, Liabeuf S, Brocard C, Haase G, Bras H, Vinay L. Activation of 5-HT_{2A} receptors upregulates the function of the neuronal K-Cl cotransporter KCC2. *Proc Natl Acad Sci U S A* 2013;110:348-353.
- [19] Braz JM, Enquist LW, Basbaum AI. Inputs to serotonergic neurons revealed by conditional viral transneuronal tracing. *J Comp Neurol* 2009;514:145-160.
- [20] Burgess SE, Gardell LR, Ossipov MH, Malan TP, Jr, Vanderah TW, Lai J, Porreca F. Time-dependent descending facilitation from the rostral ventromedial medulla maintains, but does not initiate, neuropathic pain. *J Neurosci* 2002;22:5129-5136.
- [21] Burton H, Jones EG. The posterior thalamic region and its cortical projection in New World and Old World monkeys. *J Comp Neurol* 1976;168:249-301.
- [22] Carlton SM, Leichnetz GR, Young EG, Mayer DJ. Supramedullary afferents of the nucleus raphe magnus in the rat: a study using the transcanula HRP gel and autoradiographic techniques. *J Comp Neurol* 1983;214:43-58.

- [23] Carrasquillo Y, Gereau RW 4th. Hemispheric lateralization of a molecular signal for pain modulation in the amygdala. *Mol Pain* 2008;4:24.
- [24] Chen T, Wang XL, Qu J, Wang W, Zhang T, Yanagawa Y, Wu SX, Li YQ. Neurokinin-1 receptor-expressing neurons that contain serotonin and gamma-aminobutyric acid in the rat rostroventromedial medulla are involved in pain processing. *J Pain* 2013;14:778-792.
- [25] Cho JH, Jeong MY, Choi IS, Lee HJ, Jang IS. TRPA1-like channels enhance glycinergic transmission in medullary dorsal horn neurons. *J Neurochem* 2012;122:691-701.
- [26] Coull JA, Beggs S, Boudreau D, Boivin D, Tsuda M, Inoue K, Gravel C, Salter MW, De Koninck Y. BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature* 2005;438:1017-1021.
- [27] Coull JA, Boudreau D, Bachand K, Prescott SA, Nault F, Sik A, De Koninck P, De Koninck Y. Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature* 2003;424:938-942.
- [28] Cox JJ, Reimann F, Nicholas AK, Thornton G, Roberts E, Springell K, Karbani G, Jafri H, Mannan J, Raashid Y, Al-Gazali L, Hamamy H, Valente EM, Gorman S, Williams R, McHale DP, Wood JN, Gribble FM, Woods CG. An SCN9A channelopathy causes congenital inability to experience pain. *Nature* 2006;444:894-898.
- [29] Dayas CV, Buller KM, Day TA. Neuroendocrine responses to an emotional stressor: evidence for involvement of the medial but not the central amygdala. *Eur J Neurosci* 1999;11:2312-2322.
- [30] Decosterd I, Woolf CJ. Spared nerve injury: an animal model of persistent peripheral neuropathic pain. *Pain* 2000;87:149-158.
- [31] del Camino D, Murphy S, Heiry M, Barrett LB, Earley TJ, Cook CA, Petrus MJ, Zhao M, D'Amours M, Deering N, Brenner GJ, Costigan M, Hayward NJ, Chong JA, Fanger CM, Woolf CJ, Patapoutian A, Moran MM. TRPA1 contributes to cold hypersensitivity. *J Neurosci* 2010;30:15165-15174.
- [32] Dupuis A, Wattiez AS, Pinguet J, Richard D, Libert F, Chalus M, Aissouni Y, Sion B, Ardid D, Marin P, Eschali r A, Courteix C. Increasing spinal 5-HT_{2A} receptor responsiveness mediates anti-allodynic effect and potentiates fluoxetine efficacy in neuropathic rats. Evidence for GABA release. *Pharmacol Res* 2017;118:93-103.
- [33] El Bitar N, Pollin B, Huang G, Mouraux A, Le Bars D. The rostral ventromedial medulla control of cutaneous vasomotion of paws and tail in the rat: implication for pain studies. *J Neurophysiol* 2016;115:773-789.

- [34] Eliacik K, Kanik A, Bolat N, Mertek H, Guven B, Karadas U, Dogrusoz B, Bakiler AR. Anxiety, depression, suicidal ideation, and stressful life events in non-cardiac adolescent chest pain: a comparative study about the hidden part of the iceberg. *Cardiol Young* 2016;1-6.
- [35] Emery EC, Luiz AP, Sikandar S, Magnúsdóttir R, Dong X, Wood JN. In vivo characterization of distinct modality-specific subsets of somatosensory neurons using GCaMP. *Sci Adv* 2016;2:e1600990.
- [36] England JD, Happel LT, Kline DG, Gamboni F, Thouron CL, Liu ZP, Levinson SR. Sodium channel accumulation in humans with painful neuromas. *Neurology* 1996;47:272-276.
- [37] Eslick GD, Koloski NA, Talley NJ. Sexual, physical, verbal/emotional abuse and unexplained chest pain. *Child Abuse Negl* 2011;35:601-605.
- [38] Faber ES, Sedlak P, Vidovic M, Sah P. Synaptic activation of transient receptor potential channels by metabotropic glutamate receptors in the lateral amygdala. *Neuroscience* 2006;137:781-794.
- [39] Ferretti A, Babiloni C, Gratta CD, Caulo M, Tartaro A, Bonomo L, Rossini PM, Romani GL. Functional topography of the secondary somatosensory cortex for nonpainful and painful stimuli: an fMRI study. *Neuroimage* 2003;20:1625-1638.
- [40] Fields HL, Malick A, Burstein R. Dorsal horn projection targets of ON and OFF cells in the rostral ventromedial medulla. *J Neurophysiol* 1995;74:1742-1759.
- [41] Fink KB, Gothert M. 5-HT receptor regulation of neurotransmitter release. *Pharmacol Rev* 2007;59:360-417.
- [42] Fisher HE, Brown LL, Aron A, Strong G, Mashek D. Reward, addiction, and emotion regulation systems associated with rejection in love. *J Neurophysiol* 2010;104:51-60.
- [43] Fregni F, Potvin K, Dasilva D, Wang X, Lenkinski RE, Freedman SD, Pascual-Leone A. Clinical effects and brain metabolic correlates in non-invasive cortical neuromodulation for visceral pain. *Eur J Pain* 2011;15:53-60.
- [44] Fu Y, Neugebauer V. Differential mechanisms of CRF1 and CRF2 receptor functions in the amygdala in pain-related synaptic facilitation and behavior. *J Neurosci* 2008;28:3861-3876.
- [45] Fu Y, Neugebauer V. Differential mechanisms of CRF1 and CRF2 receptor functions in the amygdala in pain-related synaptic facilitation and behavior. *J Neurosci* 2008;28:3861-3876.

- [46] Gautier A, El Ouaraki H, Bazin N, Salam S, Vodjdani G, Bourgoin S, Pezet S, Bernard JF, Hamon M. Lentiviral vector-driven inhibition of 5-HT synthesis in B3 bulbo-spinal serotonergic projections - Consequences on nociception, inflammatory and neuropathic pain in rats. *Exp Neurol* 2017;288:11-24.
- [47] Geha PY, Baliki MN, Chialvo DR, Harden RN, Paice JA, Apkarian AV. Brain activity for spontaneous pain of postherpetic neuralgia and its modulation by lidocaine patch therapy. *Pain* 2007;128:88-100.
- [48] Gjerstad J, Tjolsen A, Hole K. The effect of 5-HT_{1A} receptor stimulation on nociceptive dorsal horn neurones in rats. *Eur J Pharmacol* 1996;318:315-321.
- [49] Gojyo F, Sugiyo S, Kuroda R, Kawabata A, Varathan V, Shigenaga Y, Takemura M. Effects of somatosensory cortical stimulation on expression of c-Fos in rat medullary dorsal horn in response to formalin-induced noxious stimulation. *J Neurosci Res* 2002;68:479-488.
- [50] Gray TS. Amygdaloid CRF pathways. Role in autonomic, neuroendocrine, and behavioral responses to stress. *Ann N Y Acad Sci* 1993;697:53-60.
- [51] Greenwood-Van Meerveld B, Gibson M, Gunter W, Shepard J, Foreman R, Myers D. Stereotaxic delivery of corticosterone to the amygdala modulates colonic sensitivity in rats. *Brain Res* 2001;893:135-142.
- [52] Guliaeva NV, Levshina IP. Characteristics of free-radical oxidation and antiradical protection of the brain in adaptation to chronic stress. *Biull Eksp Biol Med* 1988;106:153-156.
- [53] Hachinski V. Headache as a presenting symptom. *Can Fam Physician* 1976;22:55-58.
- [54] Han JS, Bird GC, Neugebauer V. Enhanced group III mGluR-mediated inhibition of pain-related synaptic plasticity in the amygdala. *Neuropharmacology* 2004;46:918-926.
- [55] Hayashida K, Kimura M, Yoshizumi M, Hobo S, Obata H, Eisenach JC. Ondansetron reverses antihypersensitivity from clonidine in rats after peripheral nerve injury: role of gamma-aminobutyric acid in alpha₂-adrenoceptor and 5-HT₃ serotonin receptor analgesia. *Anesthesiology* 2012;117:389-398.
- [56] Heinricher MM, Barbaro NM, Fields HL. Putative nociceptive modulating neurons in the rostral ventromedial medulla of the rat: firing of on- and off-cells is related to nociceptive responsiveness. *Somatosens Mot Res* 1989;6:427-439.
- [57] Heinricher MM, Tavares I, Leith JL, Lumb BM. Descending control of nociception: Specificity, recruitment and plasticity. *Brain Res Rev* 2009;60:214-225.

- [58] Helmstetter FJ. Contribution of the amygdala to learning and performance of conditional fear. *Physiol Behav* 1992;51:1271-1276.
- [59] Helmstetter FJ, Tershner SA. Lesions of the periaqueductal gray and rostral ventromedial medulla disrupt antinociceptive but not cardiovascular aversive conditional responses. *J Neurosci* 1994;14:7099-7108.
- [60] Herman JP, Cullinan WE. Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. *Trends Neurosci* 1997;20:78-84.
- [61] Hokfelt T, Ljungdahl A, Steinbusch H, Verhofstad A, Nilsson G, Brodin E, Pernow B, Goldstein M. Immunohistochemical evidence of substance P-like immunoreactivity in some 5-hydroxytryptamine-containing neurons in the rat central nervous system. *Neuroscience* 1978;3:517-538.
- [62] Hossaini M, Goos JA, Kohli SK, Holstege JC. Distribution of glycine/GABA neurons in the ventromedial medulla with descending spinal projections and evidence for an ascending glycine/GABA projection. *PLoS One* 2012;7:e35293.
- [63] Hughes DJ, Scott DT, Todd AJ, Riddell JS. Lack of evidence for sprouting of Abeta afferents into the superficial laminae of the spinal cord dorsal horn after nerve section. *J Neurosci* 2003;23:9491-9499.
- [64] Hunskaar S, Hole K. The formalin test in mice: dissociation between inflammatory and non-inflammatory pain. *Pain* 1987;30:103-114.
- [65] Ikeda R, Takahashi Y, Inoue K, Kato F. NMDA receptor-independent synaptic plasticity in the central amygdala in the rat model of neuropathic pain. *Pain* 2007;127:161-172.
- [66] Itoga CA, Roltsch Hellard EA, Whitaker AM, Lu YL, Schreiber AL, Baynes BB, Baiaamonte BA, Richardson HN, Gilpin NW. Traumatic stress promotes hyperalgesia via corticotropin-releasing factor-1 receptor (CRFR1) signaling in central amygdala. *Neuropsychopharmacology* 2016;41:2463-2472.
- [67] Ji G, Fu Y, Adwanikar H, Neugebauer V. Non-pain-related CRF1 activation in the amygdala facilitates synaptic transmission and pain responses. *Mol Pain* 2013;9:2.
- [68] Ji G, Li Z, Neugebauer V. Reactive oxygen species mediate visceral pain-related amygdala plasticity and behaviors. *Pain* 2015;156:825-836.
- [69] Ji G, Neugebauer V. Pro- and anti-nociceptive effects of corticotropin-releasing factor (CRF) in central amygdala neurons are mediated through different receptors. *J Neurophysiol* 2008;99:1201-1212.

- [70] Ji G, Neugebauer V. Hemispheric lateralization of pain processing by amygdala neurons. *J Neurophysiol* 2009;102:2253-2264.
- [71] Ji G, Sun H, Fu Y, Li Z, Pais-Vieira M, Galhardo V, Neugebauer V. Cognitive impairment in pain through amygdala-driven prefrontal cortical deactivation. *J Neurosci* 2010;30:5451-5464.
- [72] Ji G, Zhang W, Mahimainathan L, Narasimhan M, Kiritoshi T, Fan X, Wang J, Green TA, Neugebauer V. 5-HT_{2C} receptor knockdown in the amygdala inhibits neuropathic-pain-related plasticity and behaviors. *J Neurosci* 2017;37:1378-1393.
- [73] Ji RR, Samad TA, Jin SX, Schmoll R, Woolf CJ. p38 MAPK activation by NGF in primary sensory neurons after inflammation increases TRPV1 levels and maintains heat hyperalgesia. *Neuron* 2002;36:57-68.
- [74] Jiang H, Fang D, Kong LY, Jin ZR, Cai J, Kang XJ, Wan Y, Xing GG. Sensitization of neurons in the central nucleus of the amygdala via the decreased GABAergic inhibition contributes to the development of neuropathic pain-related anxiety-like behaviors in rats. *Mol Brain* 2014;7:72.
- [75] Johansson O, Hokfelt T, Pernow B, Jeffcoate SL, White N, Steinbusch HW, Verhofstad AA, Emson PC, Spindel E. Immunohistochemical support for three putative transmitters in one neuron: coexistence of 5-hydroxytryptamine, substance P- and thyrotropin releasing hormone-like immunoreactivity in medullary neurons projecting to the spinal cord. *Neuroscience* 1981;6:1857-1881.
- [76] Keefe FJ, Shelby RA, Somers TJ, Varia I, Blazing M, Waters SJ, McKee D, Silva S, She L, Blumenthal JA, O'Connor J, Knowles V, Johnson P, Bradley L. Effects of coping skills training and sertraline in patients with non-cardiac chest pain: a randomized controlled study. *Pain* 2011;152:730-741.
- [77] Kerstein PC, del Camino D, Moran MM, Stucky CL. Pharmacological blockade of TRPA1 inhibits mechanical firing in nociceptors. *Mol Pain* 2009;5:19.
- [78] Kim SH, Chung JM. An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat. *Pain* 1992;50:355-363.
- [79] Kishore-Kumar R, Schafer SC, Lawlor BA, Murphy DL, Max MB. Single doses of the serotonin agonists buspirone and m-chlorophenylpiperazine do not relieve neuropathic pain. *Pain* 1989;37:223-227.
- [80] Kosugi M, Nakatsuka T, Fujita T, Kuroda Y, Kumamoto E. Activation of TRPA1 channel facilitates excitatory synaptic transmission in substantia gelatinosa neurons of the adult rat spinal cord. *J Neurosci* 2007;27:4443-4451.

- [81] Krause T, Assemer S, Taskin B, Floel A, Witte AV, Mueller K, Fiebach JB, Villringer K, Villringer A, Jungehulsing GJ. The cortical signature of central poststroke pain: gray matter decreases in somatosensory, insular, and prefrontal cortices. *Cereb Cortex* 2016;26:80-88.
- [82] Kremer M, Salvat E, Muller A, Yalcin I, Barrot M. Antidepressants and gabapentinoids in neuropathic pain: Mechanistic insights. *Neuroscience* 2016;338:183-206.
- [83] Kremeyer B, Lopera F, Cox JJ, Momin A, Rugiero F, Marsh S, Woods CG, Jones NG, Paterson KJ, Fricker FR, Villegas A, Acosta N, Pineda-Trujillo NG, Ramirez JD, Zea J, Burley MW, Bedoya G, Bennett DL, Wood JN, Ruiz-Linares A. A gain-of-function mutation in TRPA1 causes familial episodic pain syndrome. *Neuron* 2010;66:671-680.
- [84] Kronschlager MT, Drdla-Schutting R, Gassner M, Honsek SD, Teuchmann HL, Sandkuhler J. Gliogenic LTP spreads widely in nociceptive pathways. *Science* 2016;354:1144-1148.
- [85] Kulkarni B, Bentley DE, Elliott R, Julyan PJ, Boger E, Watson A, Boyle Y, El-Dereby W, Jones AK. Arthritic pain is processed in brain areas concerned with emotions and fear. *Arthritis Rheum* 2007;56:1345-1354.
- [86] Kuroda R, Kawabata A, Kawao N, Umeda W, Takemura M, Shigenaga Y. Somatosensory cortex stimulation-evoked analgesia in rats: potentiation by NO synthase inhibition. *Life Sci* 2000;66:PL271-6.
- [87] Kuroda R, Kawao N, Yoshimura H, Umeda W, Takemura M, Shigenaga Y, Kawabata A. Secondary somatosensory cortex stimulation facilitates the antinociceptive effect of the NO synthase inhibitor through suppression of spinal nociceptive neurons in the rat. *Brain Res* 2001;903:110-116.
- [88] Kwiat GC, Basbaum AI. The origin of brainstem noradrenergic and serotonergic projections to the spinal cord dorsal horn in the rat. *Somatosens Mot Res* 1992;9:157-173.
- [89] LaBuda CJ, Fuchs PN. A behavioral test paradigm to measure the aversive quality of inflammatory and neuropathic pain in rats. *Exp Neurol* 2000;163:490-494.
- [90] LeDoux JE, Iwata J, Cicchetti P, Reis DJ. Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J Neurosci* 1988;8:2517-2529.

- [91] Lee JE, Watson D, Frey-Law LA. Psychological factors predict local and referred experimental muscle pain: a cluster analysis in healthy adults. *Eur J Pain* 2013;17:903-915.
- [92] Lefaucheur JP, Drouot X, Nguyen JP. Interventional neurophysiology for pain control: duration of pain relief following repetitive transcranial magnetic stimulation of the motor cortex. *Neurophysiol Clin* 2001;31:247-252.
- [93] Lefler Y, Arzi A, Reiner K, Sukhotinsky I, Devor M. Bulbosplinal neurons of the rat rostromedial medulla are highly collateralized. *J Comp Neurol* 2008;506:960-978.
- [94] Li MJ, Liu LY, Chen L, Cai J, Wan Y, Xing GG. Chronic stress exacerbates neuropathic pain via the integration of stress-affect-related information with nociceptive information in the central nucleus of the amygdala. *Pain* 2017;158:717-739.
- [95] Li W, Neugebauer V. Block of NMDA and non-NMDA receptor activation results in reduced background and evoked activity of central amygdala neurons in a model of arthritic pain. *Pain* 2004;110:112-122.
- [96] Li W, Neugebauer V. Differential changes of group II and group III mGluR function in central amygdala neurons in a model of arthritic pain. *J Neurophysiol* 2006;96:1803-1815.
- [97] Li Z, Ji G, Neugebauer V. Mitochondrial reactive oxygen species are activated by mGluR5 through IP3 and activate ERK and PKA to increase excitability of amygdala neurons and pain behavior. *J Neurosci* 2011;31:1114-1127.
- [98] Liang M, Mouraux A, Iannetti GD. Parallel processing of nociceptive and non-nociceptive somatosensory information in the human primary and secondary somatosensory cortices: evidence from dynamic causal modeling of functional magnetic resonance imaging data. *J Neurosci* 2011;31:8976-8985.
- [99] Lindholm P, Lamusuo S, Taiminen T, Pesonen U, Lahti A, Virtanen A, Forssell H, Hietala J, Hagelberg N, Pertovaara A, Parkkola R, Jaaskelainen S. Right secondary somatosensory cortex-a promising novel target for the treatment of drug-resistant neuropathic orofacial pain with repetitive transcranial magnetic stimulation. *Pain* 2015;156:1276-1283.
- [100] Lockwood PL, Iannetti GD, Haggard P. Transcranial magnetic stimulation over human secondary somatosensory cortex disrupts perception of pain intensity. *Cortex* 2013;49:2201-2209.

- [101] Lu YF, Neugebauer V, Chen J, Li Z. Distinct contributions of reactive oxygen species in amygdala to bee venom-induced spontaneous pain-related behaviors. *Neurosci Lett* 2016;619:68-72.
- [102] Ma W, Blomqvist A, Berkley KJ. Spino-diencephalic relays through the parabrachial nucleus in the cat. *Brain Res* 1989;480:37-50.
- [103] Mains RE, Eipper BA. Coordinate synthesis of corticotropins and endorphins by mouse pituitary tumor cells. *J Biol Chem* 1978;253:651-655.
- [104] Malick A, Strassman RM, Burstein R. Trigemino-hypothalamic and reticulohypothalamic tract neurons in the upper cervical spinal cord and caudal medulla of the rat. *J Neurophysiol* 2000;84:2078-2112.
- [105] Manzano GM, Giuliano LM, Nobrega JA. A brief historical note on the classification of nerve fibers. *Arq Neuropsiquiatr* 2008;66:117-119.
- [106] Mason P. Physiological identification of pontomedullary serotonergic neurons in the rat. *J Neurophysiol* 1997;77:1087-1098.
- [107] Maxwell L, Maxwell DJ, Neilson M, Kerr R. A confocal microscopic survey of serotonergic axons in the lumbar spinal cord of the rat: co-localization with glutamate decarboxylase and neuropeptides. *Neuroscience* 1996;75:471-480.
- [108] Melzack R, Wall PD. Pain mechanisms: a new theory. *Science* 1965;150:971-979.
- [109] Merskey H, Bogduk N. Part III: Pain terms, a current list with definitions and notes on usage. In: *Classification of chronic pain. Second edition (revised)*. Seattle: IASP Press, 2011.
- [110] Migita K, Uozumi T, Arita K, Monden S. Transcranial magnetic coil stimulation of motor cortex in patients with central pain. *Neurosurgery* 1995;36:1037-9; discussion 1039-40.
- [111] Millan MJ. Descending control of pain. *Prog Neurobiol* 2002;66:355-474.
- [112] Missig G, Mei L, Vizzard MA, Braas KM, Waschek JA, Ressler KJ, Hammack SE, May V. Parabrachial pituitary adenylate cyclase-activating polypeptide activation of amygdala endosomal extracellular signal-regulated kinase signaling regulates the emotional component of pain. *Biol Psychiatry* 2017;81:671-682.
- [113] Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J* 2009;417:1-13.

- [114] Nassenstein C, Kwong K, Taylor-Clark T, Kollarik M, Macglashan DM, Braun A, Udem BJ. Expression and function of the ion channel TRPA1 in vagal afferent nerves innervating mouse lungs. *J Physiol* 2008;586:1595-1604.
- [115] Neugebauer V. Amygdala pain mechanisms. *Handb Exp Pharmacol* 2015;227:261-284.
- [116] Neugebauer V, Li W, Bird GC, Bhave G, Gereau RW, 4th. Synaptic plasticity in the amygdala in a model of arthritic pain: differential roles of metabotropic glutamate receptors 1 and 5. *J Neurosci* 2003;23:52-63.
- [117] Nitzan-Luques A, Devor M, Tal M. Genotype-selective phenotypic switch in primary afferent neurons contributes to neuropathic pain. *Pain* 2011;152:2413-2426.
- [118] Obermann M, Rodriguez-Raecke R, Naegel S, Holle D, Mueller D, Yoon MS, Theysohn N, Blex S, Diener HC, Katsarava Z. Gray matter volume reduction reflects chronic pain in trigeminal neuralgia. *Neuroimage* 2013;74:352-358.
- [119] O'Connor AB. Neuropathic pain: quality-of-life impact, costs and cost effectiveness of therapy. *Pharmacoeconomics* 2009;27:95-112.
- [120] O'Connor AB, Dworkin RH. Treatment of neuropathic pain: an overview of recent guidelines. *Am J Med* 2009;122:S22-32.
- [121] Ogawa N, Kurokawa T, Mori Y. Sensing of redox status by TRP channels. *Cell Calcium* 2016;60:115-122.
- [122] Pavlov I. Lectures on conditioned reflexes twenty-five years of objective study of the higher nervous activity (behaviour) of animals. New York, NY, USA: International Publishers, 1928.
- [123] Pedersen LH, Scheel-Kruger J, Blackburn-Munro G. Amygdala GABA-A receptor involvement in mediating sensory-discriminative and affective-motivational pain responses in a rat model of peripheral nerve injury. *Pain* 2007;127:17-26.
- [124] Pert A, Yaksh T. Localization of the antinociceptive action of morphine in primate brain. *Pharmacol Biochem Behav* 1975;3:133-138.
- [125] Peters A, Schweiger U, Pellerin L, Hubold C, Oltmanns KM, Conrad M, Schultes B, Born J, Fehm HL. The selfish brain: competition for energy resources. *Neurosci Biobehav Rev* 2004;28:143-180.
- [126] Prewitt CM, Herman JP. Hypothalamo-pituitary-adrenocortical regulation following lesions of the central nucleus of the amygdala. *Stress* 1997;1:263-280.

- [127] Price DD, Dubner R. Mechanisms of first and second pain in the peripheral and central nervous systems. *J Invest Dermatol* 1977;69:167-171.
- [128] Qin C, Greenwood-Van Meerveld B, Myers DA, Foreman RD. Corticosterone acts directly at the amygdala to alter spinal neuronal activity in response to colorectal distension. *J Neurophysiol* 2003;89:1343-1352.
- [129] Radat F, Margot-Duclot A, Attal N. Psychiatric co-morbidities in patients with chronic peripheral neuropathic pain: a multicentre cohort study. *Eur J Pain* 2013;17:1547-1557.
- [130] Raz M. The painless brain: lobotomy, psychiatry, and the treatment of chronic pain and terminal illness. *Perspect Biol Med* 2009;52:555-565.
- [131] Redgrave P, Simkins M, McHaffie JG, Stein BE. Nociceptive neurones in rat superior colliculus. II. Effects of lesions to the contralateral descending output pathway on nocifensive behaviours. *Exp Brain Res* 1996;109:197-208.
- [132] Reis FJ, Lopes D, Rodrigues J, Gosling AP, Gomes MK. Psychological distress and quality of life in leprosy patients with neuropathic pain. *Lepr Rev* 2014;85:186-193.
- [133] Ren W, Neugebauer V. Pain-related increase of excitatory transmission and decrease of inhibitory transmission in the central nucleus of the amygdala are mediated by mGluR1. *Mol Pain* 2010;6:93.
- [134] Ren W, Palazzo E, Maione S, Neugebauer V. Differential effects of mGluR7 and mGluR8 activation on pain-related synaptic activity in the amygdala. *Neuropharmacology* 2011;61:1334-1344.
- [135] Riccio A, Li Y, Moon J, Kim KS, Smith KS, Rudolph U, Gapon S, Yao GL, Tsvetkov E, Rodig SJ, Van't Veer A, Meloni EG, Carlezon WA, Jr, Bolshakov VY, Clapham DE. Essential role for TRPC5 in amygdala function and fear-related behavior. *Cell* 2009;137:761-772.
- [136] Riccio A, Li Y, Tsvetkov E, Gapon S, Yao GL, Smith KS, Engin E, Rudolph U, Bolshakov VY, Clapham DE. Decreased anxiety-like behavior and Galphaq/11-dependent responses in the amygdala of mice lacking TRPC4 channels. *J Neurosci* 2014;34:3653-3667.
- [137] Rodriguez-Raecke R, Niemeier A, Ihle K, Ruether W, May A. Brain gray matter decrease in chronic pain is the consequence and not the cause of pain. *J Neurosci* 2009;29:13746-13750.

- [138] Rodriguez-Raecke R, Niemeier A, Ihle K, Ruether W, May A. Structural brain changes in chronic pain reflect probably neither damage nor atrophy. *PLoS One* 2013;8:e54475.
- [139] Roeder Z, Chen Q, Davis S, Carlson JD, Tupone D, Heinricher MM. Parabrachial complex links pain transmission to descending pain modulation. *Pain* 2016;157:2697-2708.
- [140] Roozendaal B, Koolhaas JM, Bohus B. Central amygdala lesions affect behavioral and autonomic balance during stress in rats. *Physiol Behav* 1991;50:777-781.
- [141] Rouwette T, Vanelderen P, de Reus M, Loohuis NO, Giele J, van Egmond J, Scheenen W, Scheffer GJ, Roubos E, Vissers K, Kozicz T. Experimental neuropathy increases limbic forebrain CRF. *Eur J Pain* 2012;16:61-71.
- [142] Roychowdhury SM, Heinricher MM. Effects of iontophoretically applied serotonin on three classes of physiologically characterized putative pain modulating neurons in the rostral ventromedial medulla of lightly anesthetized rats. *Neurosci Lett* 1997;226:136-138.
- [143] Royer S, Martina M, Pare D. An inhibitory interface gates impulse traffic between the input and output stations of the amygdala. *J Neurosci* 1999;19:10575-10583.
- [144] Ruan T, Lin YJ, Hsu TH, Lu SH, Jow GM, Kou YR. Sensitization by pulmonary reactive oxygen species of rat vagal lung C-fibers: the roles of the TRPV1, TRPA1, and P2X receptors. *PLoS One* 2014;9:e91763.
- [145] Rubinstein M, Mogil JS, Japon M, Chan EC, Allen RG, Low MJ. Absence of opioid stress-induced analgesia in mice lacking beta-endorphin by site-directed mutagenesis. *Proc Natl Acad Sci U S A* 1996;93:3995-4000.
- [146] Russell WMS, Burch RL. The principles of humane experimental technique. England: Universities Federation for Animal Welfare, 1959 (as reprinted 1992).
- [147] Sawada A, Niiyama Y, Ataka K, Nagaishi K, Yamakage M, Fujimiya M. Suppression of bone marrow-derived microglia in the amygdala improves anxiety-like behavior induced by chronic partial sciatic nerve ligation in mice. *Pain* 2014;155:1762-1772.
- [148] Selye H. Stress and the General Adaptation Syndrome. *Br Med J* 1950;1:1383-1392.

- [149] Shepard JD, Barron KW, Myers DA. Corticosterone delivery to the amygdala increases corticotropin-releasing factor mRNA in the central amygdaloid nucleus and anxiety-like behavior. *Brain Res* 2000;861:288-295.
- [150] Smith DA, Flynn JP. Afferent projections related to attack sites in the pontine tegmentum. *Brain Res* 1979;164:103-119.
- [151] Sotgiu ML, Valente M, Storch R, Caramenti G, Mario Biella GE. Contribution by DRt descending facilitatory pathways to maintenance of spinal neuron sensitization in rats. *Brain Res* 2008;1188:69-75.
- [152] Spiers JG, Chen HJ, Sernia C, Lavidis NA. Activation of the hypothalamic-pituitary-adrenal stress axis induces cellular oxidative stress. *Front Neurosci* 2015;8:456.
- [153] Staaf S, Oerther S, Lucas G, Mattsson JP, Ernfors P. Differential regulation of TRP channels in a rat model of neuropathic pain. *Pain* 2009;144:187-199.
- [154] Straub RH. Insulin resistance, selfish brain, and selfish immune system: an evolutionarily positively selected program used in chronic inflammatory diseases. *Arthritis Res Ther* 2014;16 Suppl 2:S4.
- [155] Straub RH. Systemic disease sequelae in chronic inflammatory diseases and chronic psychological stress: comparison and pathophysiological model. *Ann N Y Acad Sci* 2014;1318:7-17.
- [156] Strekalova T, Spanagel R, Dolgov O, Bartsch D. Stress-induced hyperlocomotion as a confounding factor in anxiety and depression models in mice. *Behav Pharmacol* 2005;16:171-180.
- [157] Tanaka M, Matsumoto Y, Murakami T, Hisa Y, Ibata Y. The origins of catecholaminergic innervation in the rostral ventromedial medulla oblongata of the rat. *Neurosci Lett* 1996;207:53-56.
- [158] Tavares I, Lima D. The caudal ventrolateral medulla as an important inhibitory modulator of pain transmission in the spinal cord. *J Pain* 2002;3:337-346.
- [159] Taylor AM, Harris AD, Varnava A, Phillips R, Hughes O, Wilkes AR, Hall JE, Wise RG. Neural responses to a modified Stroop paradigm in patients with complex chronic musculoskeletal pain compared to matched controls: an experimental functional magnetic resonance imaging study. *BMC Psychol* 2016;4:5.
- [160] Tomoyose O, Kodama D, Ono H, Tanabe M. Presynaptic inhibitory effects of fluvoxamine, a selective serotonin reuptake inhibitor, on nociceptive excitatory synaptic transmission in spinal superficial dorsal horn neurons of adult mice. *J Pharmacol Sci* 2014;126:136-145.

- [161] Torquati K, Pizzella V, Della Penna S, Franciotti R, Babiloni C, Romani GL, Rossini PM. "Gating" effects of simultaneous peripheral electrical stimulations on human secondary somatosensory cortex: a whole-head MEG study. *Neuroimage* 2003;20:1704-1713.
- [162] Tran L, Greenwood-Van Meerveld B. Altered expression of glucocorticoid receptor and corticotropin-releasing factor in the central amygdala in response to elevated corticosterone. *Behav Brain Res* 2012;234:380-385.
- [163] Tsubokawa T, Katayama Y, Yamamoto T, Hirayama T, Koyama S. Chronic motor cortex stimulation for the treatment of central pain. *Acta Neurochir Suppl (Wien)* 1991;52:137-139.
- [164] Tsubokawa T, Katayama Y, Yamamoto T, Hirayama T, Koyama S. Chronic motor cortex stimulation in patients with thalamic pain. *J Neurosurg* 1993;78:393-401.
- [165] Uglem M, Omland PM, Engstrom M, Gravidahl GB, Linde M, Hagen K, Sand T. Non-invasive cortical modulation of experimental pain in migraine. *Clin Neurophysiol* 2016;127:2362-2369.
- [166] Ulrich-Lai YM, Xie W, Meij JT, Dolgas CM, Yu L, Herman JP. Limbic and HPA axis function in an animal model of chronic neuropathic pain. *Physiol Behav* 2006;88:67-76.
- [167] Urban MO, Smith DJ. Nuclei within the rostral ventromedial medulla mediating morphine antinociception from the periaqueductal gray. *Brain Res* 1994;652:9-16.
- [168] Valmunen T, Pertovaara A, Taiminen T, Virtanen A, Parkkola R, Jaaskelainen SK. Modulation of facial sensitivity by navigated rTMS in healthy subjects. *Pain* 2009;142:149-158.
- [169] Vanegas H, Barbaro NM, Fields HL. Tail-flick related activity in medullospinal neurons. *Brain Res* 1984;321:135-141.
- [170] Varga V, Sik A, Freund TF, Kocsis B. GABA(B) receptors in the median raphe nucleus: distribution and role in the serotonergic control of hippocampal activity. *Neuroscience* 2002;109:119-132.
- [171] Viisanen H, Ansah OB, Pertovaara A. The role of the dopamine D2 receptor in descending control of pain induced by motor cortex stimulation in the neuropathic rat. *Brain Res Bull* 2012;89:133-143.

- [172] Viisanen H, Pertovaara A. Antinociception by motor cortex stimulation in the neuropathic rat: does the locus coeruleus play a role? *Exp Brain Res* 2010;201:283-296.
- [173] Viisanen H, Pertovaara A. Roles of the rostroventromedial medulla and the spinal 5-HT(1A) receptor in descending antinociception induced by motor cortex stimulation in the neuropathic rat. *Neurosci Lett* 2010;476:133-137.
- [174] Vogt BA, Rosene DL, Pandya DN. Thalamic and cortical afferents differentiate anterior from posterior cingulate cortex in the monkey. *Science* 1979;204:205-207.
- [175] Wagman IH, Price DD. Responses of dorsal horn cells of *M. mulatta* to cutaneous and sural nerve A and C fiber stimuli. *J Neurophysiol* 1969;32:803-817.
- [176] Wagner KM, Roeder Z, Desrochers K, Buhler AV, Heinricher MM, Cleary DR. The dorsomedial hypothalamus mediates stress-induced hyperalgesia and is the source of the pronociceptive peptide cholecystokinin in the rostral ventromedial medulla. *Neuroscience* 2013;238:29-38.
- [177] Wei H, Chapman H, Saarnilehto M, Kuokkanen K, Koivisto A, Pertovaara A. Roles of cutaneous versus spinal TRPA1 channels in mechanical hypersensitivity in the diabetic or mustard oil-treated non-diabetic rat. *Neuropharmacology* 2010;58:578-584.
- [178] Wei H, Koivisto A, Saarnilehto M, Chapman H, Kuokkanen K, Hao B, Huang JL, Wang YX, Pertovaara A. Spinal transient receptor potential ankyrin 1 channel contributes to central pain hypersensitivity in various pathophysiological conditions in the rat. *Pain* 2011;152:582-591.
- [179] Wei H, Pertovaara A. 5-HT(1A) receptors in endogenous regulation of neuropathic hypersensitivity in the rat. *Eur J Pharmacol* 2006;535:157-165.
- [180] Wei H, Huang J, Hao B, Wang Y, Nian G, Ma A, Li X, Wang Y, Pertovaara A. Intrathecal administration of antioxidants attenuates mechanical pain hypersensitivity induced by REM sleep deprivation in the rat. *Scand J Pain* 2011;2:64-69.
- [181] Willis WD, Kenshalo DR, Jr, Leonard RB. The cells of origin of the primate spinothalamic tract. *J Comp Neurol* 1979;188:543-573.
- [182] Woolf CJ, Shortland P, Coggeshall RE. Peripheral nerve injury triggers central sprouting of myelinated afferents. *Nature* 1992;355:75-78.
- [183] Wu D, Huang W, Richardson PM, Priestley JV, Liu M. TRPC4 in rat dorsal root ganglion neurons is increased after nerve injury and is necessary for neurite outgrowth. *J Biol Chem* 2008;283:416-426.

[184] Xiao WH, Bennett GJ. Effects of mitochondrial poisons on the neuropathic pain produced by the chemotherapeutic agents, paclitaxel and oxaliplatin. *Pain* 2012;153:704-709.

[185] Xu Y, Day TA, Buller KM. The central amygdala modulates hypothalamic-pituitary-adrenal axis responses to systemic interleukin-1beta administration. *Neuroscience* 1999;94:175-183.

[186] You HJ, Colpaert FC, Arendt-Nielsen L. The novel analgesic and high-efficacy 5-HT_{1A} receptor agonist F 13640 inhibits nociceptive responses, wind-up, and after-discharges in spinal neurons and withdrawal reflexes. *Exp Neurol* 2005;191:174-183.

[187] Zhuo M, Gebhart GF. Biphasic modulation of spinal nociceptive transmission from the medullary raphe nuclei in the rat. *J Neurophysiol* 1997;78:746-758.